United States

Health Department Website: hhs.gov

Summary

Highlights

Drug Deaths

43%

from 6.9 to 9.9 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Number of Geriatric Providers

16%

from 31.4 to 36.4 per 100,000 adults ages 65+ between September 2019 and September 2022.

Multiple Chronic Conditions

13%

from 46% to 52% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Poverty

10%

from 9.4% to 10.3% of adults ages 65+ between 2019 and 2021.

High-speed Internet

7%

from 78.0% to 83.1% of households with adults ages 65+ between 2019 and 2021.

Food Insecurity

6%▼

from 12.6% to 11.9% of adults ages 60+ between 2019 and 2020.

Measures

U.S. Value

Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)+	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	11.8
Economic	Food Insecurity (% of adults ages 60+)	11.9%
Resources	Poverty (% of adults ages 65+)	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	\$62
and	High-speed Internet (% of households with adults ages 65+)	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	10.27
	Volunteerism (% of adults ages 65+)	22.1%
	·	22.1%
Physical Enviro		
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	7.8
Quality	Drinking Water Violations (% of community water systems)	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	32.3%
Clinical Care*		
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	96.1%
Quality of Care	Hospice Care (% of Medicare decedents)	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	1,482
Behaviors*		
Nutrition	Exercise (% of adults ages 65+)	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	26.09
Tobacco Use	Smoking (% of adults ages 65+)	8.9%
Health Outcon	nes*	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	9.9
	Excessive Drinking (% of adults ages 65+)	7.0%
	Frequent Mental Distress (% of adults ages 65+)	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	2,151
•	Early Death Racial Disparity (ratio)***	1.5
	Falls (% of adults ages 65+)	27.1%
Physical Health		
Physical Health	Frequent Physical Distress (% of adults ages 65+)	15.8%
Physical Health	Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	15.8% 52%
Physical Health	Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74) Obesity (% of adults ages 65+)	15.8% 52% 29.5%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

** Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of excessive drinking
- High level of community support expenditures
- · Low prevalence of severe housing problems

Challenges

- · High early death rate
- · Low percentage of households with high-speed internet
- · Low home health care worker rate

Highlights

Physical Inactivity

from 33.7% to 41.3% of adults ages 65+ in fair or better health between 2018 and 2021.

Food Insecurity

from 19.3% to 15.3% of adults ages 60+ between 2015-2016 and 2019-2020.

Multiple Chronic Conditions

from 58% to 64% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	44	-0.672	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	44	20.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	38	15.8	11.8
Economic	Food Insecurity (% of adults ages 60+)	44	15.3%	11.9%
Resources	Poverty (% of adults ages 65+)	42	11.4%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	29	62.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	7	\$104	\$62
ind	High-speed Internet (% of households with adults ages 65+)	47	76.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	26	15.5%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	44	81	_
	Volunteerism (% of adults ages 65+)	47	16.9%	22.1%
Physical Enviro	onment*	8	0.543	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	28	7.8	7.8
Quality	Drinking Water Violations (% of community water systems)	30	0.3%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	3	24.0%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	7	24.6%	32.3%
Clinical Care*		30	-0.051	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	18	2.7%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	20	36.2	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	48	22.2	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	12	79.2%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	27	67.0%	67.5%
services	Pneumonia Vaccination (% of adults ages 65+)	33	70.1%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	9	97.0%	96.19
Care	Hospice Care (% of Medicare decedents)	28	46.6%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	20	36.3%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	46	2,033	1,482
Behaviors*		49	-1.474	
Nutrition	Exercise (% of adults ages 65+)	45	15.1%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	37	5.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	49	41.3%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	48	30.3%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	44	12.3%	8.9%
Health Outcon	nes*	45	-0.677	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	43	9.4%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	2	5.5	9.9
	Excessive Drinking (% of adults ages 65+)	6	4.7%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	29	8.3%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	31	18.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	47	2,928	2,151
,	Early Death Racial Disparity (ratio)***	_	1.3	1.5
Physical Health	Falls (% of adults ages 65+)	22	28.0%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	43	19.2%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	50	64%	52%
	Obesity (% of adults ages 65+)	25	30.9%	29.59
	Teeth Extractions (% of adults ages 65+)	44	18.7%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.



Summary

Strengths

- · Low prevalence of multiple chronic conditions
- High level of community support expenditures
- · Low poverty rate

Challenges

- · High prevalence of excessive drinking
- · Low percentage of older adults with a dedicated health care provider
- · High suicide rate

Highlights

Early Deaths

from 1,667 to 2,163 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Home Health Care Workers

from 75.4 to 65.6 aides per 1,000 adults ages 65+ between 2019 and 2021.

High-speed Internet

from 79.8% to 86.2% of households with adults ages 65+ between 2017 and 2021.

Measure	9 S	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	6	0.660	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	47	23.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	20	11.5	11.8
Economic	Food Insecurity (% of adults ages 60+)	29	12.1%	11.9%
Resources	Poverty (% of adults ages 65+)	9	8.5%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	16	88.2	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	2	\$164	\$62
and	High-speed Internet (% of households with adults ages 65+)	8	86.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	_	_	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	21	47	_
	Volunteerism (% of adults ages 65+)	19	26.8%	22.1%
Physical Enviro		43	-0.293	22.170
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	15	7.0	7.8
Quality	Drinking Water Violations (% of community water systems)	47	4.8%	0.8%
Housing	Housing Cost Burden (% of households with adults	11	25.5%	31.8%
nousing	ages 65+)‡	11		31.070
	Severe Housing Problems (% of small households with adults ages 62+)	28	29.9%	32.3%
Clinical Care*		45	-0.688	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	41	4.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	45	21.5	36.4
	Home Health Care Workers (workers per 1,000 adults	9	65.6	60.3
	ages 65+)		00.0	00.0
Preventive	Cancer Screenings (% of adults ages 65-75)	40	74.0%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	49	58.5%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	45	64.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	49	90.9%	96.1%
Care	Hospice Care (% of Medicare decedents)	49	30.1%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	2	52.2%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	10	1,072	1,482
Behaviors*	Medicare perienciaries ages 05-74)	27	0.020	
Nutrition	Exercise (% of adults ages 65+)	4	29.0%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	7	9.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or	25	30.9%	31.2%
	better health)	20	30.376	01.270
Sleep Health	Insufficient Sleep (% of adults ages 65+)	40	28.3%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	41	11.8%	8.9%
Health Outcon	nes*	34	-0.236	
	Cognitive Difficulty (% of adults ages 65+)‡	4	6.0%	7.8%
	cognitive billiouity (70 of addite ages 60)			9.9
	Drug Deaths (deaths per 100,000 adults ages 65+)‡	22	9.8	0.0
		22 48	9.8 10.5%	7.0%
	Drug Deaths (deaths per 100,000 adults ages 65+)‡			7.0%
	Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	48	10.5%	7.0% 8.5%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)† Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	48 19	10.5% 7.6%	7.0%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	48 19 44	10.5% 7.6% 24.9	7.0% 8.5% 16.9
Health Mortality	Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***	48 19 44 29 —	10.5% 7.6% 24.9 2,163 3.8	7.0% 8.5% 16.9 2,151 1.5
Health Mortality	Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+)	48 19 44 29 — 50	10.5% 7.6% 24.9 2,163 3.8 36.5%	7.0% 8.5% 16.9 2,151 1.5 27.1%
Health Mortality	Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	48 19 44 29 —	10.5% 7.6% 24.9 2,163 3.8	7.0% 8.5% 16.9 2,151
Behavioral Health Mortality Physical Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	48 19 44 29 — 50 24 2	10.5% 7.6% 24.9 2,163 3.8 36.5% 15.5% 36%	7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8% 52%
Health Mortality	Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	48 19 44 29 — 50 24	10.5% 7.6% 24.9 2,163 3.8 36.5% 15.5%	7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.



Summary

Strengths

- · High percentage of households with high-speed internet
- · Low risk of social isolation
- · High hospice care use

Challenges

- · High suicide rate
- · Low percentage of older adults with a dedicated health care provider
- Low prevalence of fruit and vegetable consumption

Highlights

Number of Geriatric Providers

from 24.4 to 27.9 per 100,000 adults ages 65+ between September 2019 and September 2022.

Early Deaths

from 1,962 to 2,220 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

Poverty

from 8.6% to 9.6% of adults ages 65+ between 2018 and 2021.

Measure	es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	27	0.163	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	43	20.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	26	13.1	11.8
Economic	Food Insecurity (% of adults ages 60+)	32	12.8%	11.9%
Resources	Poverty (% of adults ages 65+)	28	9.6%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	30	61.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	31	\$36	\$62
and	High-speed Internet (% of households with adults ages 65+)	6	87.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	16	11.9%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	7	33	_
	Volunteerism (% of adults ages 65+)	37	19.5%	22.1%
Physical Enviro	onment*	45	-0.380	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	49	10.5	7.8
Quality	Drinking Water Violations (% of community water systems)	34	0.5%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	28	29.0%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	29	30.0%	32.3%
Clinical Care*		32	-0.167	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	28	3.2%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	33	27.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	14	50.1	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	43	72.6%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	44	63.1%	67.5%
Jei vices	Pneumonia Vaccination (% of adults ages 65+)	37	69.1%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	42	93.5%	96.1%
Care	Hospice Care (% of Medicare decedents)	4	54.7%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	22	35.6%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	8	1,020	1,482
Behaviors*	· ·	19	0.156	
Nutrition	Exercise (% of adults ages 65+)	9	27.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	40	4.6%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	11	27.4%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	20	24.1%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	35	9.4%	8.9%
Health Outcon	nes*	31	-0.150	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	24	7.5%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	32	10.5	9.9
	Excessive Drinking (% of adults ages 65+)	37	8.0%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	36	8.8%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	43	24.5	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	31	2,220	2,151
	Early Death Racial Disparity (ratio)***		1.7	1.5
Physical Health	Falls (% of adults ages 65+)	17	27.0%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	31	16.3%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	21	49%	52%
	Obesity (% of adults ages 65+)	17	28.3%	29.5%
	Teeth Extractions (% of adults ages 65+)	28	13.3%	13.4%
Overall		30	-0.010	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- · Low prevalence of excessive drinking
- High geriatric provider rate
- · Low prevalence of severe housing problems

Challenges

- · High early death rate
- Low percentage of households with high-speed internet
- · High prevalence of physical inactivity

Highlights

Home Health Care Workers

from 45.2 to 32.7 aides per 1,000 adults ages 65+ between 2017 and 2021.

Obesity

from 27.6% to 33.4% of adults ages 65+ between 2016 and 2021.

High-speed Internet

from 70.4% to 76.6% of households with adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	45	-0.806	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	36	18.7	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	47	18.6	11.8
Economic	Food Insecurity (% of adults ages 60+)	28	11.5%	11.9%
Resources	Poverty (% of adults ages 65+)	44	11.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	49	37.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	18	\$52	\$62
and .	High-speed Internet (% of households with adults ages 65+)	46	76.6%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	39	18.0%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	43	79	_
	Volunteerism (% of adults ages 65+)	39	18.7%	22.1%
Physical Enviro	onment*	14	0.487	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	37	8.3	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	12	25.6%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	4	24.0%	32.3%
Clinical Care*		35	-0.200	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	32	3.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	7	48.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	37	32.7	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	44	72.2%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	39	64.4%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	27	70.7%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	32	95.7%	96.1%
Care	Hospice Care (% of Medicare decedents)	32	45.7%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	19	37.2%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	40	1,767	1,482
Behaviors*		41	-0.728	
Nutrition	Exercise (% of adults ages 65+)	38	19.8%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	19	7.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	46	40.0%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	18	23.5%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	47	13.0%	8.9%
Health Outcon	nes*	48	-0.949	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	48	9.8%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	6	6.1	9.9
	Excessive Drinking (% of adults ages 65+)	7	4.8%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	42	9.7%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	34	19.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	45	2,875	2,151
•	Early Death Racial Disparity (ratio)***	_	1.2	1.5
Physical Health	Falls (% of adults ages 65+)	47	32.9%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	47	20.8%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	34	55%	52%
	Obesity (% of adults ages 65+)	41	33.4%	29.5%
	Teeth Extractions (% of adults ages 65+)	48	21.5%	13.4%
	. ,	44	-0.640	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · High percentage of households with high-speed internet
- · High home health care worker rate
- · Low prevalence of physical inactivity

Challenges

- · High prevalence of excessive drinking
- · Low volunteerism rate
- · High prevalence of severe housing problems

Highlights

Volunteerism

from 22.9% to 15.2% of adults ages 65+ between 2019 and 2021.

Food Insecurity

from 16.3% to 11.4% of adults ages 60+ between 2012-2013 and 2019-2020.

Early Deaths

from 1,452 to 1,823 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	36	-0.208	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	11	9.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	9	9.0	11.8
Economic	Food Insecurity (% of adults ages 60+)	27	11.4%	11.99
Resources	Poverty (% of adults ages 65+)	40	11.1%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	37	54.5	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	34	\$35	\$62
and .	High-speed Internet (% of households with adults ages 65+)	4	87.6%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	17	12.1%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	37	63	_
	Volunteerism (% of adults ages 65+)	49	15.2%	22.19
Physical Enviro	onment*	49	-1.140	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	50	12.8	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	50	40.5%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	49	42.1%	32.3
Clinical Care*		33	-0.177	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	32	3.4%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	33	27.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	2	120.4	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	49	65.5%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	40	64.0%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	44	65.4%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	27	96.1%	96.1
Care	Hospice Care (% of Medicare decedents)	41	42.3%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	12	40.3%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	16	1,195	1,48
Behaviors*		11	0.382	
Nutrition	Exercise (% of adults ages 65+)	31	21.3%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	9	8.7%	7.39
Activity	Physical Inactivity (% of adults ages 65+ in fair or	6	25.4%	31.29
Sleep Health	better health)	34	26.79/	26.0
Tobacco Use	Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	6	26.7% 7.3%	26.0 8.99
				0.97
Health Outcon		20	0.215	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	38	8.8%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	42	13.0	9.9
	Excessive Drinking (% of adults ages 65+)	46	9.9%	7.09
	Frequent Mental Distress (% of adults ages 65+)	44	10.1%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	14	15.5	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	12	1,823	2,15
m	Early Death Racial Disparity (ratio)***		1.7	1.5
Physical Health	Falls (% of adults ages 65+)	15	26.5%	27.19
	Frequent Physical Distress (% of adults ages 65+)	35	16.4%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	13	46%	529
	Obesity (% of adults ages 65+)	3	23.9%	29.5
	Teeth Extractions (% of adults ages 65+)	3	9.0%	13.4
Overall		34	-0.025	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of multiple chronic conditions
- · Low poverty rate
- Low prevalence of physical inactivity

Challenges

- High suicide rate
- Low level of community support expenditures
- Low home health care worker rate

Highlights

Drug Deaths

70%

from 7.0 to 11.9 deaths per 100,000 adults ages 65+ between 2014-2016 and 2019-2021.

Number of Geriatric Providers

18%

from 36.8 to 43.3 per 100,000 adults ages 65+ between September 2019 and September 2022.

High-speed Internet

13%

from 78.0% to 88.1% of households with adults ages 65+ between 2016 and 2021.

Measure	28	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	9	0.489	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	35	18.6	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	15	10.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	25	11.0%	11.9%
Resources	Poverty (% of adults ages 65+)	4	7.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	23	76.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	23	\$43	\$62
and	High-speed Internet (% of households with adults ages 65+)	2	88.1%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	46	25.0%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	5	29	_
	Volunteerism (% of adults ages 65+)	8	31.9%	22.1%
Physical Enviro		26	0.213	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	19	7.3	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	37	32.3%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	35	32.3%	32.3%
Clinical Care*	,	5	0.693	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	7	2.3%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	10	43.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	21	45.5	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	30	76.6%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	11	72.0%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	1	76.3%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	30	96.0%	96.1%
Care	Hospice Care (% of Medicare decedents)	14	50.4%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	16	38.3%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	2	772	1,482
Behaviors*		2	1.040	
Nutrition	Exercise (% of adults ages 65+)	2	29.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	19	7.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	1	21.4%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	3	20.1%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	23	8.5%	8.9%
Health Outcon	nes*	4	0.612	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	10	6.6%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	39	11.9	9.9
	Excessive Drinking (% of adults ages 65+)	20	6.7%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	10	7.1%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	45	25.8	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	9	1,745	2,151
•	Early Death Racial Disparity (ratio)***	_	1.8	1.5
Physical Health	Falls (% of adults ages 65+)	36	29.8%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	10	13.1%	15.8%
Physical Health	Multiple Chronic Conditions (% of Medicare beneficiaries			52%
		5	40%	0270
	ages 65-74)			
		5 2 6	40% 22.9% 9.6%	29.5% 13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.
Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.
Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Connecticut

Overall Rank

Summary

Strengths

- · Low early death rate
- Low prevalence of frequent physical distress
- · Low poverty rate

Challenges

- · Low level of community support expenditures
- · Low hospice care use
- · High prevalence of severe housing problems

Highlights

Drug Deaths

from 7.1 to 12.6 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Multiple Chronic Conditions

from 45% to 51% of Medicare beneficiaries ages 65+ between 2020 and 2021.

High-speed Internet

from 79.8% to 85.2% of households with adults ages 65+ between 2019 and 2021.

<i>Measure</i>	2 S	State Rank	State Value	U.S Valu
Social & Econo	omic Factors*	17	0.309	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	6	6.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	5	7.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	33	12.9%	11.99
Resources	Poverty (% of adults ages 65+)	11	8.6%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	36	\$32	\$62
ind	High-speed Internet (% of households with adults ages 65+)	13	85.2%	83.19
Ingagement	Low-care Nursing Home Residents (% of residents)	31	16.5%	15.2
	Risk of Social Isolation (index from 1-100, adults ages 65+)	23	50	_
	Volunteerism (% of adults ages 65+)	16	28.4%	22.19
Physical Enviro	onment*	32	0.077	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	8	6.1	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	46	37.7%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	46	38.7%	32.3
Clinical Care*		7	0.555	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	12	2.5%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	4	57.6	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	11	58.6	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	16	78.8%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	13	71.6%	67.5
services	Pneumonia Vaccination (% of adults ages 65+)	16	72.9%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	17	96.5%	96.1
Care	Hospice Care (% of Medicare decedents)	39	43.8%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	20	36.3%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	22	1,375	1,48
Behaviors*		7	0.742	
Nutrition	Exercise (% of adults ages 65+)	9	27.3%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	5	10.2%	7.39
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	21	29.8%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	26	25.4%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	3	6.5%	8.99
Health Outcon	nes*	2	0.889	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	15	6.9%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	41	12.6	9.9
	Excessive Drinking (% of adults ages 65+)	27	7.1%	7.09
	Frequent Mental Distress (% of adults ages 65+)	10	7.1%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	11	14.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	2	1,570	2,15
	Early Death Racial Disparity (ratio)***		1.5	1.5
Physical Health	Falls (% of adults ages 65+)	2	21.3%	27.19
	Frequent Physical Distress (% of adults ages 65+)	1	10.6%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	23	51%	52%
	•			
	Obesity (% of adults ages 65+) Teeth Extractions (% of adults ages 65+)	10	26.6%	29.5

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low suicide rate
- · Low poverty rate
- · High volunteerism rate

Challenges

- · High prevalence of multiple chronic conditions
- High prevalence of obesity
- · High prevalence of physical inactivity

Highlights

Drug Deaths

from 5.9 to 10.0 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Food Insecurity

from 12.3% to 9.3% of adults ages 60+ between 2014-2015 and 2019-2020.

Flu Vaccination Rates

from 63.4% to 73.7% of adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	2	0.902	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	8	8.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	36	15.2	11.8
Economic	Food Insecurity (% of adults ages 60+)	10	9.3%	11.9%
Resources	Poverty (% of adults ages 65+)	2	6.9%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	22	77.4	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	24	\$42	\$62
and .	High-speed Internet (% of households with adults ages 65+)	7	86.5%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	22	13.6%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	2	19	_
	Volunteerism (% of adults ages 65+)	6	34.2%	22.1%
Physical Enviro	onment*	23	0.243	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	39	8.4	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	12	25.6%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	22	28.2%	32.3%
Clinical Care*		16	0.269	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	32	3.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	26	30.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	29	41.0	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	25	77.5%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	5	73.7%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	18	72.8%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	37	95.2%	96.1%
Care	Hospice Care (% of Medicare decedents)	12	51.9%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	6	45.5%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	34	1,607	1,482
Behaviors*		33	-0.048	
Nutrition	Exercise (% of adults ages 65+)	18	24.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	16	7.6%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or	33	33.2%	31.2%
Sleep Health	better health) Insufficient Sleep (% of adults ages 65+)	39	28.0%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	23	8.5%	8.9%
Health Outcon		27	-0.021	0.576
	Cognitive Difficulty (% of adults ages 65+)*		5.9%	7.8%
Behavioral Health		3		
	Drug Deaths (deaths per 100,000 adults ages 65+)‡	23	10.0	9.9
	Excessive Drinking (% of adults ages 65+)	44 15	8.9%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	15	7.3%	8.5%
Mantality:	Suicide (deaths per 100,000 adults ages 65+)	5	11.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***	20	1,959 1.2	2,151 1.5
Physical Health	Falls (% of adults ages 65+)	10		27.1%
- iiysicai neaith			25.6%	
	Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	15	13.9%	15.8%
	ages 65-74)	48	60%	52%
	Obesity (% of adults ages 65+)	37	32.9%	29.5%
	Teeth Extractions (% of adults ages 65+)	23	12.9%	13.4%
	<u> </u>			

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- · High level of community support expenditures
- · High hospice care use
- High percentage of households with high-speed internet

Challenges

- High prevalence of multiple chronic conditions
- · Low home health care worker rate
- · Low volunteerism rate

Highlights

Early Deaths

from 1,598 to 2,027 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Number of Geriatric Providers

from 33.5 to 37.2 per 100,000 adults ages 65+ between September 2021 and September 2022.

Poverty

from 10.2% to 11.0% of adults ages 65+ between 2017 and 2021.

Measure	2 S	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	19	0.291	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	25	14.7	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	29	13.5	11.8
Economic	Food Insecurity (% of adults ages 60+)	35	13.4%	11.9%
Resources	Poverty (% of adults ages 65+)	38	11.0%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	3	\$154	\$62
and	High-speed Internet (% of households with adults ages 65+)	11	86.0%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	14	11.3%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	23	50	_
	Volunteerism (% of adults ages 65+)	45	17.1%	22.1%
Physical Envir		35	-0.037	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	19	7.3	7.8
Quality	Drinking Water Violations (% of community water systems)	40	1.3%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	41	33.7%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	36	33.4%	32.3%
Clinical Care*		34	-0.181	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	_	_	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	17	37.2	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	50	14.6	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	5	80.4%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	_	_	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	_		69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	_	_	96.1%
Care	Hospice Care (% of Medicare decedents)	2	56.2%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	15	38.9%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	33	1,585	1,482
Behaviors*		21	0.134	
Nutrition	Exercise (% of adults ages 65+)	8	27.8%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	_	_	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	_	_	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	28	25.5%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	_	_	8.9%
Health Outcor	nes*	29	-0.100	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	28	7.7%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	16	8.3	9.9
	Excessive Drinking (% of adults ages 65+)	_	_	7.0%
	Frequent Mental Distress (% of adults ages 65+)	_	_	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	35	19.9	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	24	2,027	2,151
	Early Death Racial Disparity (ratio)**‡	_	1.4	1.5
Physical Health	Falls (% of adults ages 65+)	4	24.2%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	_	_	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	44	58%	52%
	Obesity (% of adults ages 65+)	_	_	29.5%
	Teeth Extractions (% of adults ages 65+)	31	13.9%	13.4%
Overall		25	0.051	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

Note: Florida data are missing for 2021 Behavioral Risk Factor Surveillance System measures due to the state's exclusion from the Centers for Disease Control and Prevention's annual aggregate data set.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

 $For measure \ descriptions, source \ details \ and \ methodology, \ visit \ Americas Health Rankings. org.$

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.

Georgia

State Health Department Website: dph.georgia.gov

Summary

Strengths

- · Low prevalence of excessive drinking
- · Low prevalence of obesity
- · High hospice care use

Challenges

- · Low level of community support expenditures
- · High prevalence of food insecurity
- · Low flu vaccination rate

Highlights

Early Deaths

from 2,015 to 2,550 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Home Health Care Workers

from 28.4 to 22.9 aides per 1,000 adults ages 65+ between 2018 and 2021.

High-speed Internet

from 75.9% to 82.8% of households with adults ages 65+ between 2019 and 2021. **Overall Rank**

<i>Measure</i>	2 S	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	39	-0.281	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	33	16.1	13.1
amily Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	44	17.3	11.8
conomic	Food Insecurity (% of adults ages 60+)	42	14.3%	11.9%
Resources	Poverty (% of adults ages 65+)	31	10.0%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	20	79.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	49	\$18	\$62
ind	High-speed Internet (% of households with adults ages 65+)	25	82.8%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	23	14.1%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	34	61	_
	Volunteerism (% of adults ages 65+)	44	17.4%	22.19
Physical Enviro	onment*	30	0.123	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	33	8.2	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	24	28.1%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	31	30.4%	32.39
Clinical Care*	adults ages 02-7	43	-0.485	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	44	4.8%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	21	34.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	47	22.9	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	33	76.0%	75.99
Clinical	Flu Vaccination (% of adults ages 65+)	42	63.2%	67.59
Services	Pneumonia Vaccination (% of adults ages 65+)	32	70.2%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	17	96.5%	96.19
Care	Hospice Care (% of Medicare decedents)	18	48.3%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	44	25.2%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	39	1,738	1,482
Behaviors*	,	31	-0.044	
Nutrition	Exercise (% of adults ages 65+)	23	23.4%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	15	7.7%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	35	33.5%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	33	26.5%	26.09
obacco Use	Smoking (% of adults ages 65+)	27	8.6%	8.9%
Health Outcon		35	-0.249	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	39	8.9%	7.8%
-lealth	Drug Deaths (deaths per 100,000 adults ages 65+)*	13	7.4	9.9
	Excessive Drinking (% of adults ages 65+)	18	6.6%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	32	8.4%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	25	17.6	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	42	2,550	2,15
•	Early Death Racial Disparity (ratio)***	_	1.2	1.5
hysical Health	Falls (% of adults ages 65+)	18	27.2%	27.19
	Frequent Physical Distress (% of adults ages 65+)	29	15.9%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	34	55%	52%
	Obesity (% of adults ages 65+)	18	28.7%	29.59
	Teeth Extractions (% of adults ages 65+)	41	17.5%	13.49
			.,	.5. 17

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

Ties in value are assigned equal ranks.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- · Low early death rate
- · Low prevalence of obesity
- · Low prevalence of physical inactivity

Challenges

- · Low volunteerism rate
- · Low home health care worker rate
- · High prevalence of severe housing problems

Highlights

Home Health Care Workers

from 32.8 to 23.4 aides per 1,000 adults ages 65+ between 2020 and 2021.

Food Insecurity

from 10.2% to 7.5% of adults ages 60+ between 2017-2018 and 2019-2020.

Multiple Chronic Conditions

from 41% to 47% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	7	0.633	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	5	4.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	1	5.8	11.8
Economic	Food Insecurity (% of adults ages 60+)	5	7.5%	11.9%
Resources	Poverty (% of adults ages 65+)	22	9.4%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	16	\$54	\$62
and	High-speed Internet (% of households with adults ages 65+)	9	86.1%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	1	3.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	14	42	_
	Volunteerism (% of adults ages 65+)	41	18.4%	22.1%
Physical Enviro		12	0.503	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	1	3.8	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	41	33.7%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	44	37.0%	32.3%
Clinical Care*		15	0.322	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	26	3.0%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	8	46.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	46	23.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	5	80.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	35	65.6%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	49	60.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	4	97.4%	96.1%
Care	Hospice Care (% of Medicare decedents)	22	48.0%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	1	73.3%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	1	736	1,482
Behaviors*		25	0.042	
Nutrition	Exercise (% of adults ages 65+)	15	25.8%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	28	6.4%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	6	25.4%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	50	38.6%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	4	7.0%	8.9%
Health Outcon	nes*	1	1.132	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	40	9.0%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	46	15.3	9.9
	Excessive Drinking (% of adults ages 65+)	34	7.8%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	10	7.1%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	9	13.8	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	1	1,552	2,151
•	Early Death Racial Disparity (ratio)***	_	1.5	1.5
Physical Health	Falls (% of adults ages 65+)	2	21.3%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	3	11.9%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	16	47%	52%
	Obesity (% of adults ages 65+)	1	16.5%	29.5%
	Teeth Extractions (% of adults ages 65+)	1	5.6%	13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- Low prevalence of frequent mental distress
- · Low risk of social isolation
- · Low prevalence of physical inactivity

Challenges

- · High suicide rate
- · Low geriatric provider rate
- · Low flu vaccination rate

Highlights

Multiple Chronic Conditions

from 35% to 42% of Medicare beneficiaries ages 65+ between 2020 and 2021.

High-speed Internet

from 73.8% to 84.3% of households with adults ages 65+ between 2016 and 2021.

Early Deaths

from 1,748 to 1,971 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

earm Deaths (deaths per 100,000 adults ages 65+)‡ earm Deaths (deaths per 100,000 adults ages 65+)‡ old Insecurity (% of adults ages 60+) verty (% of adults ages 60+) verty (% of adults ages 65+) IAP Reach (participants per 100 adults ages 60+ n poverty) mmunity Support Expenditures (dollars per adult ages 60+) gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) ment* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) pusing Cost Burden (% of households with adults ges 65+)‡ vere Housing Problems (% of small households with dults ages 62+) orided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+) ment Health Care Workers (workers per 1,000 adults	21 41 18 17 25 38 48 18 37 4 9 21 16 42 19	0.247 19.7 10.8 9.7% 9.5% 54.4 \$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	13.1 11.8 11.9% 10.3% 81.0 \$62 83.1% 15.2% - 22.1% 7.8 0.8% 31.8% 32.3%
otor Vehicle Deaths (deaths per 100,000 adults ages 65+)* od Insecurity (% of adults ages 60+) verty (% of adults ages 65+) IAP Reach (participants per 100 adults ages 60+) poverty) IAP Reach (participants per 100 adults ages 60+) poverty) IAP Reach (participants per 100 adults ages 60+) poverty) IAP Reach (participants per 100 adults ages 60+) poverty) IAP Reach (participants per 100 adults ages 60+) IAP Reach (participants per 100 adults ages 60+) IAP Reach (participants per 100,000 adults ages 60+) IAP Reach (participants per 100,000 adults ages 65+) IAP Reach (participants ages 60+) IAP Reach (participants age	18 17 25 38 48 18 37 4 9 21 16 42 19 18	10.8 9.7% 9.5% 54.4 \$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	11.8 11.9% 10.3% 81.0 \$62 83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
od Insecurity (% of adults ages 60+) verty (% of adults ages 65+) IAP Reach (participants per 100 adults ages 60+ n poverty) mmunity Support Expenditures (dollars per adult ages 60+) gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) susing Cost Burden (% of households with adults ges 65+) vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	17 25 38 48 18 37 4 9 21 16 42 19 18	9.7% 9.5% 54.4 \$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	11.9% 10.3% 81.0 \$62 83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
verty (% of adults ages 65+) IAP Reach (participants per 100 adults ages 60+ n poverty) IAP Reach (participants per 100 adults ages 60+ n poverty) IAP Reach (participants per 100 adults ages 60+ n poverty) IAP Reach (participants per 100 adults ages 60+) IAP Reach (participants) IAP Reach (parti	25 38 48 18 37 4 9 21 16 42 19 18	9.5% 54.4 \$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	10.3% 81.0 \$62 83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
AP Reach (participants per 100 adults ages 60+ n poverty) mmunity Support Expenditures (dollars per adult ages 60+) gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) using Cost Burden (% of households with adults ges 65+) vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	38 48 18 37 4 9 21 16 42 19 18	\$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	\$1.0 \$62 83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
n poverty) mmunity Support Expenditures (dollars per adult ages 60+) gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) using Cost Burden (% of households with adults ges 65+) vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	48 18 37 4 9 21 16 42 19 18	\$19 84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	\$62 83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
mmunity Support Expenditures (dollars per adult ages 60+) gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) susing Cost Burden (% of households with adults ges 65+) vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	18 37 4 9 21 16 42 19 18	84.3% 17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	83.1% 15.2% — 22.1% 7.8 0.8% 31.8%
gh-speed Internet (% of households with adults ages 65+) w-care Nursing Home Residents (% of residents) sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) susing Cost Burden (% of households with adults ges 65+)+ vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	37 4 9 21 16 42 19 18	17.4% 25 31.1% 0.307 7.1 1.8% 27.1%	15.2% — 22.1% 7.8 0.8% 31.8%
sk of Social Isolation (index from 1-100, adults ages 65+) lunteerism (% of adults ages 65+) nent* Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) susing Cost Burden (% of households with adults ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	4 9 21 16 42 19 18	25 31.1% 0.307 7.1 1.8% 27.1%	7.8 0.8% 31.89
Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) susing Cost Burden (% of households with adults ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	9 21 16 42 19 18	31.1% 0.307 7.1 1.8% 27.1%	7.8 0.8% 31.8%
Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) busing Cost Burden (% of households with adults ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) oriatric Providers (providers per 100,000 adults ages 65+)	21 16 42 19 18 24	7.1 1.8% 27.1%	7.8 0.8% 31.8%
Pollution (micrograms of fine particles per cubic meter) inking Water Violations (% of community water systems) busing Cost Burden (% of households with adults ges 65+)‡ vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) viriatric Providers (providers per 100,000 adults ages 65+)	16 42 19 18 24	7.1 1.8% 27.1% 27.1%	0.8% 31.8%
inking Water Violations (% of community water systems) busing Cost Burden (% of households with adults ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) briatric Providers (providers per 100,000 adults ages 65+)	19 18 24	1.8% 27.1% 27.1%	0.8% 31.8%
ousing Cost Burden (% of households with adults ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) viatric Providers (providers per 100,000 adults ages 65+)	19 18 24	27.1% 27.1%	31.8%
ges 65+)* vere Housing Problems (% of small households with dults ages 62+) oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	18 24	27.1%	
oided Care Due to Cost (% of adults ages 65+) eriatric Providers (providers per 100,000 adults ages 65+)	24		32.39
eriatric Providers (providers per 100,000 adults ages 65+)			
eriatric Providers (providers per 100,000 adults ages 65+)		0.112	
	18	2.7%	3.4%
me Health Care Workers (workers per 1,000 adults	50	16.8	36.4
ges 65+)	12	55.9	60.3
ncer Screenings (% of adults ages 65-75)	31	76.5%	75.99
ı Vaccination (% of adults ages 65+)	45	62.5%	67.5%
eumonia Vaccination (% of adults ages 65+)	40	68.3%	69.79
dicated Health Care Provider (% of adults ages 65+)	39	94.7%	96.19
ospice Care (% of Medicare decedents)	7	54.1%	46.79
rsing Home Quality (% of beds rated four or five stars)	4	47.5%	32.69
eventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	3	794	1,482
	22	0.122	
ercise (% of adults ages 65+)	41	19.4%	23.1%
uit and Vegetable Consumption (% of adults ages 65+)	41	4.5%	7.3%
ysical Inactivity (% of adults ages 65+ in fair or etter health)	10	26.7%	31.29
sufficient Sleep (% of adults ages 65+)	4	20.4%	26.09
noking (% of adults ages 65+)	23	8.5%	8.9%
*	13	0.353	
ognitive Difficulty (% of adults ages 65+)*	42	9.3%	7.8%
ug Deaths (deaths per 100,000 adults ages 65+)‡	27	10.2	9.9
cessive Drinking (% of adults ages 65+)	39	8.2%	7.0%
equent Mental Distress (% of adults ages 65+)	6	7.0%	8.5%
icide (deaths per 100,000 adults ages 65+)	42	23.4	16.9
rly Death (deaths per 100,000 adults ages 65-74)	22	1,971	2,15
rly Death Racial Disparity (ratio)**‡	_	1.1	1.5
lls (% of adults ages 65+)	12	26.0%	27.1%
equent Physical Distress (% of adults ages 65+)	17	14.4%	15.89
ultiple Chronic Conditions (% of Medicare beneficiaries ges 65-74)	7	42%	52%
pesity (% of adults ages 65+)	16	27.7%	29.5%
coity (70 or addite ages co 1)	19	12.0%	13.49
	ait and Vegetable Consumption (% of adults ages 65+) ysical Inactivity (% of adults ages 65+ in fair or etter health) sufficient Sleep (% of adults ages 65+) noking (% of adults ages 65+) gnitive Difficulty (% of adults ages 65+) gug Deaths (deaths per 100,000 adults ages 65+) equent Mental Distress (% of adults ages 65+) sicide (deaths per 100,000 adults ages 65+) rly Death (deaths per 100,000 adults ages 65-74) rly Death Racial Disparity (ratio)*** Is (% of adults ages 65+) equent Physical Distress (% of adults ages 65+) ultiple Chronic Conditions (% of Medicare beneficiaries	### Procise (% of adults ages 65+) ### and Vegetable Consumption (% of adults ages 65+) ### ages 65+ in fair or ### ages 65+ i	### Provise (% of adults ages 65+) ### 19.4% #### 19.4% ####################################

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.



Summary

Strengths

- · Low prevalence of excessive drinking
- High flu vaccination rate
- · Low prevalence of smoking

Challenges

- · High prevalence of obesity
- · High prevalence of food insecurity
- · High prevalence of physical inactivity

Highlights

High-speed Internet

from 68.9% to 81.5% of households with adults ages 65+ between 2016 and 2021.

Poverty

from 8.6% to 10.0% of adults ages 65+ between 2019 and 2021.

Early Deaths

from 2,149 to 2,039 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

Measure	2S	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	33	-0.108	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	7	8.6	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	13	10.3	11.8
Economic	Food Insecurity (% of adults ages 60+)	38	13.7%	11.9%
Resources	Poverty (% of adults ages 65+)	31	10.0%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	13	98.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	28	\$39	\$62
and	High-speed Internet (% of households with adults ages 65+)	30	81.5%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	24	14.8%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	39	64	_
	Volunteerism (% of adults ages 65+)	29	22.9%	22.1%
Physical Enviro		46	-0.423	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	48	9.8	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults	39	32.5%	31.8%
	ages 65+)+		52.070	51.070
	Severe Housing Problems (% of small households with adults ages 62+)	38	34.1%	32.3%
Clinical Care*	addits ages 021)	38	-0.280	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	28	3.2%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	29	30.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	17	48.9	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	48	67.2%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	9	72.9%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	25	70.9%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	15	96.6%	96.1%
Care	Hospice Care (% of Medicare decedents)	26	47.1%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	48	22.3%	32.6%
	Preventable Hospitalizations (discharges per 100,000			
	Medicare beneficiaries ages 65-74)	43	1,821	1,482
Behaviors*		28	-0.014	
Nutrition	Exercise (% of adults ages 65+)	5	28.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	42	4.4%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	41	36.0%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	36	26.8%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	2	6.2%	8.9%
Health Outcon	nes*	10	0.399	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	17	7.0%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	23	10.0	9.9
	Excessive Drinking (% of adults ages 65+)	9	5.2%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	14	7.2%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	6	12.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	25	2,039	2,151
	Early Death Racial Disparity (ratio)***	_	1.6	1.5
Physical Health	Falls (% of adults ages 65+)	1	20.0%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	31	16.3%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries	23	51%	52%
		20	0170	
	ages 65-74)			29.5%
		34 33	32.4% 14.9%	29.5% 13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



State State

115

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.



Summary

Strengths

- Low prevalence of frequent mental distress
- · Low poverty rate
- · Low prevalence of avoiding care due to cost

Challenges

- · High early death rate
- High prevalence of obesity
- · High prevalence of physical inactivity

Highlights

Number of Geriatric Providers

from 30.9 to 40.1 per 100,000 adults ages 65+ between September 2020 and September 2022.

Food Insecurity

from 13.5% to 9.8% of adults ages 60+ between 2017-2018 and 2019-2020.

Multiple Chronic Conditions

from 48% to 54% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	23	0.217	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	31	16.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	27	13.3	11.8
Economic	Food Insecurity (% of adults ages 60+)	19	9.8%	11.9%
Resources	Poverty (% of adults ages 65+)	11	8.6%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	42	49.5	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	34	\$35	\$62
nd	High-speed Internet (% of households with adults ages 65+)	32	80.9%	83.1%
ngagement	Low-care Nursing Home Residents (% of residents)	9	10.5%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	26	51	_
	Volunteerism (% of adults ages 65+)	15	29.6%	22.1%
Physical Enviro		28	0.187	22.17
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	46	9.0	7.8
Quality	Drinking Water Violations (% of community water systems)	33	0.4%	0.8%
		7		31.89
Housing	Housing Cost Burden (% of households with adults ages 65+)*		24.9%	
	Severe Housing Problems (% of small households with adults ages 62+)	13	26.0%	32.39
Clinical Care*		27	-0.012	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	12	2.5%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	14	40.1	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	34	34.2	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	37	74.5%	75.99
Clinical	Flu Vaccination (% of adults ages 65+)	30	66.6%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	15	73.1%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	23	96.3%	96.19
Care	Hospice Care (% of Medicare decedents)	25	47.3%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	32	31.5%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	41	1,792	1,482
Behaviors*	,	38	-0.222	
Nutrition	Exercise (% of adults ages 65+)	27	21.8%	23.1%
nd Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	13	8.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	39	35.2%	31.29
	Insufficient Sleep (% of adults ages 65+)	20	24.1%	26.09
Sleep Health				8.9%
•	Smoking (% of adults ages 65+)	37	11.0%	
obacco Use	Smoking (% of adults ages 65+)	37 38	11.0%	
Tobacco Use Health Outcon	nes*	38	-0.299	
Tobacco Use Health Outcon Behavioral	nes* Cognitive Difficulty (% of adults ages 65+)‡	38	-0.299 7.4%	7.8%
obacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)*	38 22 18	-0.299 7.4% 8.7	7.8% 9.9
obacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+)	22 18 12	7.4% 8.7 6.0%	7.8% 9.9 7.0%
obacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	22 18 12 10	7.4% 8.7 6.0% 7.1%	7.8% 9.9 7.0% 8.5%
obacco Use Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	22 18 12 10 24	7.4% 8.7 6.0% 7.1% 17.4	7.8% 9.9 7.0% 8.5% 16.9
obacco Use Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	22 18 12 10 24 41	7.4% 8.7 6.0% 7.1% 17.4 2,514	7.8% 9.9 7.0% 8.5% 16.9 2,151
obacco Use Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***‡	22 18 12 10 24 41	7.4% 8.7 6.0% 7.1% 17.4 2.514 1.3	7.8% 9.9 7.0% 8.5% 16.9 2,15'
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+)	22 18 12 10 24 41 - 40	7.4% 8.7 6.0% 7.1% 17.4 2.514 1.3 30.2%	7.8% 9.9 7.0% 8.5% 16.9 2,15' 1.5
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***‡	22 18 12 10 24 41 - 40 27	7.4% 8.7 6.0% 7.1% 17.4 2.514 1.3 30.2% 15.7%	7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5 27.1%
Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality Physical Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+)	22 18 12 10 24 41 - 40	7.4% 8.7 6.0% 7.1% 17.4 2.514 1.3 30.2%	7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5 27.1%
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	22 18 12 10 24 41 - 40 27	7.4% 8.7 6.0% 7.1% 17.4 2.514 1.3 30.2% 15.7%	7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8%
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	22 18 12 10 24 41 - 40 27 30	7.4% 8.7 6.0% 71% 17.4 2.514 1.3 30.2% 15.7% 54%	7.8% 9.9 7.0% 8.5% 16.9 2,151

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.



Summary

Strengths

- · Low prevalence of frequent mental distress
- Low prevalence of food insecurity
- · Low prevalence of avoiding care due to cost

Challenges

- · High prevalence of obesity
- Low percentage of households with high-speed internet
- · Low geriatric provider rate

Highlights

Drug Deaths

from 3.1 to 5.8 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Poverty

from 6.6% to 8.4% of adults ages 65+ between 2017 and 2021.

Number of Geriatric Providers

from 18.6 to 22.1 per 100,000 adults ages 65+ between September 2019 and September 2022.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	30	0.042	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	18	13.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	19	11.0	11.8
Economic	Food Insecurity (% of adults ages 60+)	7	8.5%	11.9%
Resources	Poverty (% of adults ages 65+)	8	8.4%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	44	47.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	22	\$44	\$62
and	High-speed Internet (% of households with adults ages 65+)	40	79.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	42	20.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	8	35	_
	Volunteerism (% of adults ages 65+)	24	24.8%	22.1%
Physical Enviro	· · · · · · · · · · · · · · · · · · ·	7	0.593	22.170
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	28	7.8	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults	15	25.9%	31.8%
Judiniy	ages 65+) [‡]	10	20.070	01.070
	Severe Housing Problems (% of small households with adults ages 62+)	5	24.5%	32.3%
Clinical Care*		10	0.478	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	3	1.9%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	41	22.1	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	27	41.7	60.3
	Cancer Screenings (% of adults ages 65-75)	21	78.2%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	7	73.2%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	3	75.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	36	95.3%	96.1%
Care	Hospice Care (% of Medicare decedents)	10	52.7%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	18	38.0%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	13	1,156	1,482
Behaviors*	•	31	-0.044	
Nutrition	Exercise (% of adults ages 65+)	31	21.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	31	5.9%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	31	33.0%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	10	22.7%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	21	8.4%	8.9%
Health Outcon	nes*	23	0.120	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	7	6.1%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	4	5.8	9.9
	Excessive Drinking (% of adults ages 65+)	29	7.3%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	5	6.9%	8.5%
		20	16.4	16.9
	Suicide (deaths per 100.000 adults ades 65±1			2,151
Mortality	Suicide (deaths per 100,000 adults ages 65+) Farly Death (deaths per 100,000 adults ages 65-74)		2.054	
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	26	2,054 2.0	
	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡	26 —	2.0	1.5
	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+)	26 - 36	2.0	1.5 27.1%
	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	26 —	2.0	
	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	26 — 36 9 21	2.0 29.8% 12.7% 49%	1.5 27.1% 15.8% 52%
Mortality Physical Health	Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	26 — 36 9	2.0 29.8% 12.7%	1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Summary

Strengths

- · Low prevalence of frequent mental distress
- · High volunteerism rate
- · High hospice care use

Challenges

- · High prevalence of obesity
- Low geriatric provider rate
- · High prevalence of physical inactivity

Highlights

Poverty

from 7.2% to 9.1% of adults ages 65+ between 2019 and 2021.

Community Support Expenditures

from \$33 to \$41 per adult ages 60+ between 2020 and 2021.

Multiple Chronic Conditions

from 44% to 52% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Measure	2 S	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	31	-0.044	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	30	15.9	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	48	19.5	11.8
Economic	Food Insecurity (% of adults ages 60+)	26	11.2%	11.9%
Resources	Poverty (% of adults ages 65+)	16	9.1%	10.3%
	SNAP Reach (participants per 100 adults ages 60+	45	46.4	81.0
	in poverty)			
Social Support	Community Support Expenditures (dollars per adult ages 60+)	27	\$41	\$62
ınd	High-speed Internet (% of households with adults ages 65+)	28	82.4%	83.1%
Ingagement	Low-care Nursing Home Residents (% of residents)	47	26.0%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	21	47	- 10.27
	Volunteerism (% of adults ages 65+)	5	34.9%	22.19
Physical Enviro		33	0.050	22.17
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	40	8.5	7.8
Quality				
	Drinking Water Violations (% of community water systems)	41	1.7%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)*	32	29.9%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	15	26.8%	32.39
Clinical Care*		19	0.217	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	14	2.6%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	40	24.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	18	47.9	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	38	74.4%	75.99
Clinical	Flu Vaccination (% of adults ages 65+)	18	70.1%	67.59
	Pneumonia Vaccination (% of adults ages 65+)	11	74.4%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	17	96.5%	96.19
Care	Hospice Care (% of Medicare decedents)	11	52.1%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	28	32.9%	32.69
		20	32.970	32.07
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	21	1,358	1,482
Behaviors*	•	34	-0.116	
Nutrition	Exercise (% of adults ages 65+)	30	21.5%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	34	5.7%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	30	32.5%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	12	23.1%	26.09
obacco Use	Smoking (% of adults ages 65+)	30	9.1%	8.9%
lealth Outcon		24	0.112	0.57
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	24	7.5%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	_		
		9	7.0	9.9
	Excessive Drinking (% of adults ages 65+)	12	6.0%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	6	7.0%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	26	17.8	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	33	2,277	2,15
	Early Death Racial Disparity (ratio)***	_	1.6	1.5
hysical Health	Falls (% of adults ages 65+)	33	29.4%	27.19
	Frequent Physical Distress (% of adults ages 65+)	13	13.5%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	26	52%	52%
	•			00.50
	Obesity (% of adults ages 65+)	28	311%	745
	Obesity (% of adults ages 65+) Teeth Extractions (% of adults ages 65+)	28 21	31.1% 12.4%	29.59 13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- Low prevalence of excessive drinking
- High prevalence of fruit and vegetable consumption
- Low prevalence of severe housing problems

Challenges

- High prevalence of frequent mental distress
- High risk of social isolation
- High prevalence of physical inactivity

Highlights

Drug Deaths

77%

from 6.1 to 10.8 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Food Insecurity

39%

from 14.4% to 20.0% of adults ages 60+ between 2017-2018 and 2019-2020.

High-speed Internet

9%

from 71.8% to 78.3% of households with adults ages 65+ between 2019 and 2021.

Measure	es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	48	-0.962	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	37	19.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	46	17.9	11.8
Economic	Food Insecurity (% of adults ages 60+)	48	20.0%	11.9%
Resources	Poverty (% of adults ages 65+)	44	11.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	43	49.4	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	37	\$31	\$62
and	High-speed Internet (% of households with adults ages 65+)	44	78.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	19	12.9%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	46	85	_
	Volunteerism (% of adults ages 65+)	46	17.0%	22.1%
Physical Enviro		17	0.437	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	37	8.3	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	6	24.8%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	10	25.3%	32.3%
Clinical Care*		47	-0.740	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	45	5.0%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	38	25.1	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	42	27.5	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	18	78.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	38	64.5%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	42	65.9%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	21	96.4%	96.1%
Care	Hospice Care (% of Medicare decedents)	43	42.1%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	38	29.1%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	48	2,104	1,482
Behaviors*		46	-1.192	
Nutrition	Exercise (% of adults ages 65+)	49	10.5%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	11	8.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	43	37.2%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	49	32.8%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	41	11.8%	8.9%
Health Outcon	nes*	49	-1.197	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	49	10.2%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	36	10.8	9.9
	Excessive Drinking (% of adults ages 65+)	11	5.7%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	48	12.2%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	33	19.6	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	46	2,906	2,151
	Early Death Racial Disparity (ratio)***	_	1.2	1.5
Physical Health	Falls (% of adults ages 65+)	41	30.6%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	48	21.9%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	44	58%	52%
	Obesity (% of adults ages 65+)	45	35.3%	29.5%
	Teeth Extractions (% of adults ages 65+)	50	22.4%	13.4%
Overall		48	-0.952	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.
Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.
Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low suicide rate
- · Low prevalence of excessive drinking
- High percentage of older adults with a dedicated health care provider

Challenges

- · High prevalence of food insecurity
- · High prevalence of physical inactivity
- · High prevalence of avoiding care due to cost

Highlights

Physical Inactivity

from 31.5% to 40.5% of adults ages 65+ in fair or better health between 2017 and 2021.

Poverty

from 12.4% to 14.1% of adults ages 65+ between 2018 and 2021.

High-speed Internet

from 71.1% to 75.5% of households with adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	49	-1.336	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	29	15.3	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	41	16.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	50	21.0%	11.9%
Resources	Poverty (% of adults ages 65+)	50	14.1%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	36	55.4	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	19	\$51	\$62
and	High-speed Internet (% of households with adults ages 65+)	48	75.5%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	31	16.5%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	49	99	_
	Volunteerism (% of adults ages 65+)	50	14.7%	22.1%
Physical Enviro		20	0.323	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	23	7.4	7.8
Quality	Drinking Water Violations (% of community water systems)	42	1.8%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	21	27.6%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	12	25.8%	32.39
Clinical Care*		46	-0.728	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	47	5.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	47	19.4	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	19	46.6	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	18	78.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	42	63.2%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	35	69.5%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	17	96.5%	96.1%
Care	Hospice Care (% of Medicare decedents)	27	46.7%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	50	19.7%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	49	2,176	1,482
Behaviors*		48	-1.366	
Nutrition	Exercise (% of adults ages 65+)	43	18.6%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	44	4.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	47	40.5%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	46	29.7%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	45	12.5%	8.9%
Health Outcon		47	-0.771	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	47	9.7%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	28	10.3	9.9
	Excessive Drinking (% of adults ages 65+)	16	6.3%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	45	10.2%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	13	15.1	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	43	2,721	2,151
•	Early Death Racial Disparity (ratio)***	_	1.3	1.5
Physical Health	Falls (% of adults ages 65+)	28	28.5%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	45	19.7%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	44	58%	52%
	Obesity (% of adults ages 65+)	45	35.3%	29.59
	Teeth Extractions (% of adults ages 65+)	40	17.2%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of multiple chronic conditions
- High volunteerism rate
- · Low prevalence of avoiding care due to cost

Challenges

- · High suicide rate
- · High prevalence of excessive drinking
- · High prevalence of physical inactivity

Highlights

Food Insecurity

from 15.5% to 13.1% of adults ages 60+ between 2016-2017 and 2019-2020.

Home Health Care Workers

from 60.2 to 53.5 aides per 1,000 adults ages 65+ between 2018 and 2021.

High-speed Internet

from 76.6% to 84.2% of households with adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	11	0.471	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	20	14.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	23	11.9	11.8
Economic	Food Insecurity (% of adults ages 60+)	34	13.1%	11.9%
Resources	Poverty (% of adults ages 65+)	19	9.2%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	17	85.5	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	28	\$39	\$62
and	High-speed Internet (% of households with adults ages 65+)	19	84.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	2	4.4%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	19	46	_
	Volunteerism (% of adults ages 65+)	12	29.9%	22.1%
Physical Enviro	· · · · · · · · · · · · · · · · · · ·	4	0.820	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	6	5.1	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults	33	30.3%	31.8%
	ages 65+)*	23	22.070	01.070
	Severe Housing Problems (% of small households with adults ages 62+)	30	30.1%	32.3%
Clinical Care*	Ŭ ,	4	0.714	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	3	1.9%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	29	30.3	36.4
	Home Health Care Workers (workers per 1,000 adults	13	53.5	60.3
	ages 65+)	10	00.0	00.0
	Cancer Screenings (% of adults ages 65-75)	3	81.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	13	71.6%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	19	72.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	6	97.2%	96.1%
Care	Hospice Care (% of Medicare decedents)	9	52.9%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	9	42.8%	32.6%
	Preventable Hospitalizations (discharges per 100,000	12	1,145	1,482
	Medicare beneficiaries ages 65-74)			
Rehaviors*	medical of Schollance ages so 7 17	39	-0 270	
		39	-0.270	23.1%
Nutrition	Exercise (% of adults ages 65+)	40	19.5%	23.1%
Nutrition and Physical	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+)	40 29	19.5% 6.3%	7.3%
Nutrition and Physical	Exercise (% of adults ages 65+)	40	19.5%	7.3%
Nutrition and Physical Activity	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or	40 29	19.5% 6.3%	7.3% 31.2%
Nutrition and Physical Activity Sleep Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health)	40 29 38	19.5% 6.3% 35.1%	
Nutrition and Physical Activity Sleep Health Tobacco Use	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	40 29 38	19.5% 6.3% 35.1% 25.6%	7.3% 31.2% 26.0%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes*	40 29 38 29 12	19.5% 6.3% 35.1% 25.6% 7.8%	7.3% 31.2% 26.0%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	40 29 38 29 12 22	19.5% 6.3% 35.1% 25.6% 7.8% 0.150	7.3% 31.2% 26.0% 8.9%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+)	40 29 38 29 12 22 20	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2%	7.3% 31.2% 26.0% 8.9%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+)	40 29 38 29 12 22 20 8	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100.000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	40 29 38 29 12 22 20 8 39	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	40 29 38 29 12 22 20 8 39 19 37	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) **Thestar Cognitive Difficulty (% of adults ages 65+) **Drug Deaths (deaths per 100,000 adults ages 65+) **Excessive Drinking (% of adults ages 65+) **Frequent Mental Distress (% of adults ages 65+) **Suicide (deaths per 100,000 adults ages 65+) **Early Death (deaths per 100,000 adults ages 65-74)	40 29 38 29 12 22 20 8 39 19	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) **Cognitive Difficulty (% of adults ages 65+) **Drug Deaths (deaths per 100,000 adults ages 65+) **Excessive Drinking (% of adults ages 65+) **Frequent Mental Distress (% of adults ages 65+) **Suicide (deaths per 100,000 adults ages 65+) **Early Death (deaths per 100,000 adults ages 65-74) **Early Death Racial Disparity (ratio)***	40 29 38 29 12 22 20 8 39 19 37 16	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1 1.901	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+)	40 29 38 29 12 22 20 8 39 19 37 16 —	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1 1.901 — 29.1%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5
Behaviors* Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality Physical Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) Smoking (% of adults ages 65+) Thest Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100.000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100.000 adults ages 65+) Early Death (deaths per 100.000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	40 29 38 29 12 22 20 8 39 19 37 16	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1 1.901	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2,151
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) Smoking (% of adults ages 65+) Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	40 29 38 29 12 22 20 8 39 19 37 16 — 31 15	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1 1.901 — 29.1% 13.9% 43%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5 27.1% 15.8% 52%
Nutrition and Physical Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) Smoking (% of adults ages 65+) Thest Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100.000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100.000 adults ages 65+) Early Death (deaths per 100.000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	40 29 38 29 12 22 20 8 39 19 37 16 —	19.5% 6.3% 35.1% 25.6% 7.8% 0.150 7.2% 6.7 8.2% 7.6% 20.1 1.901 — 29.1% 13.9%	7.3% 31.2% 26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Maryland

State Health Department Website: health.maryland.gov/Pages/Home.aspx

Overall Rank

Summary

Strengths

- Low prevalence of frequent physical distress
- · Low poverty rate
- · High pneumonia vaccination rate

Challenges

- · High prevalence of multiple chronic conditions
- · Low level of community support expenditures
- · Low hospice care use

Highlights

Volunteerism

from 38.2% to 22.7% of adults ages 65+ between 2017 and 2021.

Food Insecurity

from 11.6% to 9.5% of adults ages 60+ between 2017-2018 and 2019-2020.

Home Health Care Workers

from 33.1 to 28.0 aides per 1,000 adults ages 65+ between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	mic Factors*	13	0.430	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	10	9.7	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	8	8.7	11.8
Economic	Food Insecurity (% of adults ages 60+)	12	9.5%	11.9%
Resources	Poverty (% of adults ages 65+)	16	9.1%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	40	\$30	\$62
and	High-speed Internet (% of households with adults ages 65+)	15	85.1%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	5	8.8%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	23	50	
	Volunteerism (% of adults ages 65+)	30	22.7%	22.1%
Physical Enviro		24	0.240	22.170
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	13	6.7	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults	35	31.8%	31.8%
nousing	ages 65+)‡			
	Severe Housing Problems (% of small households with adults ages 62+)	38	34.1%	32.3%
Clinical Care*		11	0.465	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	14	2.6%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	5	49.2	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	41	28.0	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	10	79.5%	75.9%
OI: 1	Flu Vaccination (% of adults ages 65+)	10	72.8%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	11	74.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	2	97.9%	96.1%
Care	Hospice Care (% of Medicare decedents)	38	44.4%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	27	33.1%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	29	1,484	1,482
Behaviors*		12	0.372	
Nutrition	Exercise (% of adults ages 65+)	21	24.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	14	7.8%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	15	28.2%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	24	25.3%	26.0%
		0	7.6%	8.9%
•	Smoking (% of adults ages 65+)	9		
Tobacco Use	Smoking (% of adults ages 65+) 1es*	7	0.452	
Tobacco Use Health Outcon	nes*		0.452	7.8%
Tobacco Use Health Outcon Behavioral	nes* Cognitive Difficulty (% of adults ages 65+)‡	7	0.452 6.2%	7.8% 9.9
Tobacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)*	7 8 48	0.452 6.2% 16.6	9.9
Tobacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+)	8 48 16	0.452 6.2% 16.6 6.3%	9.9 7.0%
Tobacco Use Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	8 48 16 19	6.2% 16.6 6.3% 7.6%	9.9 7.0% 8.5%
Tobacco Use Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	8 48 16 19 8	6.2% 16.6 6.3% 7.6% 13.5	9.9 7.0% 8.5% 16.9
Tobacco Use Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	8 48 16 19	6.2% 16.6 6.3% 7.6% 13.5 1,934	9.9 7.0% 8.5% 16.9 2,151
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡	8 48 16 19 8 18	0.452 6.2% 16.6 6.3% 7.6% 13.5 1.934 1.4	9.9 7.0% 8.5% 16.9 2,151 1.5
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+)	8 48 16 19 8 18 —	0.452 6.2% 16.6 6.3% 7.6% 13.5 1.934 1.4 24.6%	9.9 7.0% 8.5% 16.9 2,151 1.5 27.1%
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	8 48 16 19 8 18	0.452 6.2% 16.6 6.3% 7.6% 13.5 1.934 1.4	9.9 7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8%
Tobacco Use Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	8 48 16 19 8 18 5 10	0.452 6.2% 16.6 6.3% 7.6% 13.5 1.934 1.4 24.6% 13.1% 55%	9.9 7.0% 8.5% 16.9 2.151 1.5 27.1% 15.8%
Tobacco Use Health Outcon Behavioral Health Mortality Physical Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	8 48 16 19 8 18 - 5 10	0.452 6.2% 16.6 6.3% 7.6% 13.5 1.934 1.4 24.6% 13.1%	9.9 7.0% 8.5% 16.9 2,151 1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Massachusetts

Overall Rank

Summary

Strengths

- · Low early death rate
- · High level of community support expenditures
- · High home health care worker rate

Challenges

- · High prevalence of excessive drinking
- · High risk of social isolation
- · High prevalence of severe housing problems

Highlights

Drug Deaths

from 6.4 to 10.0 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Poverty

from 9.1% to 10.6% of adults ages 65+ between 2019 and 2021.

Number of Geriatric Providers

from 59.3 to 63.6 per 100,000 adults ages 65+ between September 2019 and September 2022.

<i>A</i> easure	2 S	State Rank	State Value	U.S Valu
Social & Econo	omic Factors*	12	0.462	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	1	3.5	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	3	6.8	11.8
conomic	Food Insecurity (% of adults ages 60+)	12	9.5%	11.99
Resources	Poverty (% of adults ages 65+)	35	35 10.6%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	1	\$272	\$62
ınd	High-speed Internet (% of households with adults ages 65+)	13	85.2%	83.1
ngagement	Low-care Nursing Home Residents (% of residents)	13	11.2%	15.2
	Risk of Social Isolation (index from 1-100, adults ages 65+)	36	62	_
	Volunteerism (% of adults ages 65+)	34	21.6%	22.19
Physical Enviro	onment*	41	-0.173	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	19	7.3	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)*	47	38.6%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	47	38.9%	32.3
Clinical Care*		3	0.812	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	22	2.8%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	2	63.6	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	4	88.8	60.
al: : I	Cancer Screenings (% of adults ages 65-75)	1	84.2%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	4	74.7%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	23	71.2%	69.7
Quality of Care	Dedicated Health Care Provider (% of adults ages 65+)	3	97.8%	96.1
Jare	Hospice Care (% of Medicare decedents)	35	45.1%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	23	35.4%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	38	1,729	1,48
Behaviors*		29	-0.034	
Nutrition	Exercise (% of adults ages 65+)	38	19.8%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	21	7.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	14	27.8%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	41	28.5%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	9	7.6%	8.99
Health Outcon		6	0.534	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	28	7.7%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	23	10.0	9.9
	Excessive Drinking (% of adults ages 65+)	41	8.5%	7.09
	Frequent Mental Distress (% of adults ages 65+)	25	7.9%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	4	9.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	3	1,618	2,15
	Early Death Racial Disparity (ratio)***	_	1.2	1.5
Physical Health	Falls (% of adults ages 65+)	9	25.2%	27.19
	Frequent Physical Distress (% of adults ages 65+)	12	13.2%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	23	51%	529
	Obesity (% of adults ages 65+)	8	26.3%	29.5
	Teeth Extractions (% of adults ages 65+)			13.4

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of food insecurity
- · High volunteerism rate
- · High prevalence of fruit and vegetable consumption

Challenges

- · High prevalence of excessive drinking
- · High prevalence of obesity
- · Low level of community support expenditures

Highlights

Drug Deaths

from 8.5 to 10.6 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

High-speed Internet

from 69.8% to 82.3% of households with adults ages 65+ between 2016 and 2021.

Hospice Care

from 53.7% to 49.8% of Medicare decedents between 2017 and 2020.

Measure	es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	20	0.285	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	17	12.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	24	12.6	11.8
Economic	Food Insecurity (% of adults ages 60+)	10	9.3%	11.9%
Resources	Poverty (% of adults ages 65+)	20	9.3%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	19	80.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	31	\$36	\$62
and	High-speed Internet (% of households with adults ages 65+)	29	82.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	18	12.3%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	31	58	_
	Volunteerism (% of adults ages 65+)	10	30.9%	22.1%
Physical Enviro	·	22	0.247	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	33	8.2	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	23	28.0%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	24	28.6%	32.3%
Clinical Care*		18	0.223	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	24	2.9%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	14	40.1	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	24	43.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	17	78.5%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	22	69.8%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	20	72.0%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	15	96.6%	96.1%
Care	Hospice Care (% of Medicare decedents)	16	49.8%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	17	38.2%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	45	1,878	1,482
Behaviors*		30	-0.042	
Nutrition	Exercise (% of adults ages 65+)	25	23.2%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	9	8.7%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	23	30.2%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	31	26.3%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	37	11.0%	8.9%
Health Outcon	nes*	32	-0.176	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	24	7.5%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	34	10.6	9.9
	Excessive Drinking (% of adults ages 65+)	34	7.8%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	25	7.9%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	17	16.1	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	32	2,274	2,151
•	Early Death Racial Disparity (ratio)***	_	1.6	1.5
Physical Health	Falls (% of adults ages 65+)	32	29.3%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	25	15.6%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	30	54%	52%
	Obesity (% of adults ages 65+)	29	31.4%	29.5%
	Teeth Extractions (% of adults ages 65+)	26	13.0%	13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of frequent mental distress
- Low prevalence of food insecurity
- High home health care worker rate

Challenges

- · High prevalence of falls
- Low level of community support expenditures
- Low percentage of older adults with a dedicated health care provider

Highlights

Drug Deaths

73%

from 3.7 to 6.4 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Excessive Drinking

26%

from 9.2% to 6.8% of adults ages 65+ between 2016 and 2021.

Poverty

15%

from 7.4% to 8.5% of adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	14	0.413	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	9	9.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	5	7.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	3	6.6%	11.9%
Resources	Poverty (% of adults ages 65+)	9	8.5%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	34	57.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	37	\$31	\$62
and .	High-speed Internet (% of households with adults ages 65+)	24	83.0%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	34	17.1%	15.2
	Risk of Social Isolation (index from 1-100, adults ages 65+)	6	32	_
	Volunteerism (% of adults ages 65+)	11	30.7%	22.19
Physical Enviro	onment*	27	0.207	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	23	7.4	7.8
Quality	Drinking Water Violations (% of community water systems)	39	1.1%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)*	29	29.4%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	27	29.5%	32.3
Clinical Care*		1	0.975	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	7	2.3%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	3	62.6	36.
	Home Health Care Workers (workers per 1,000 adults ages 65+)	3	113.9	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	8	80.0%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	12	71.7%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	24	71.0%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	38	95.1%	96.1
Care	Hospice Care (% of Medicare decedents)	8	53.6%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	7	43.8%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	15	1,185	1,48
Behaviors*		8	0.648	
Nutrition	Exercise (% of adults ages 65+)	13	26.7%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	22	7.0%	7.39
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	13	27.6%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	2	20.0%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	21	8.4%	8.99
Health Outcon		3	0.714	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	9	6.3%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	7	6.4	9.9
	Excessive Drinking (% of adults ages 65+)	21	6.8%	7.09
	Frequent Mental Distress (% of adults ages 65+)	2	5.9%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	11	14.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	5	1,694	2,15
	Early Death Racial Disparity (ratio)***		2.6	1.5
Physical Health	Falls (% of adults ages 65+)	30	28.6%	27.19
	Frequent Physical Distress (% of adults ages 65+)	4	12.0%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	11	44%	52%
	Obesity (% of adults ages 65+)	24	30.8%	29.5
	Teeth Extractions (% of adults ages 65+)	2	8.8%	13.4
Overall		4	0.637	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Mississippi **Overall Rank** State Health Department Website: msdh.ms.g

Summary

Strengths

- · Low prevalence of frequent mental distress
- · Low prevalence of excessive drinking
- · Low prevalence of severe housing problems

Challenges

- · High early death rate
- · High poverty rate
- · High prevalence of food insecurity

Highlights

Home Health Care Workers

from 27.8 to 41.7 aides per 1,000 adults ages 65+ between 2018 and 2021.

Food Insecurity

from 17.7% to 20.7% of adults ages 60+ between 2016-2017 and 2019-2020.

High-speed Internet

from 63.8% to 72.5% of households with adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	50	-1.357	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	37	19.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	49	20.2	11.8
Economic	Food Insecurity (% of adults ages 60+)	49	20.7%	11.9%
Resources	Poverty (% of adults ages 65+)	49	13.8%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	41	50.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	42	\$29	\$62
and .	High-speed Internet (% of households with adults ages 65+)	50	72.5%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	34	17.1%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	50	100	_
	Volunteerism (% of adults ages 65+)	39	18.7%	22.19
Physical Enviro	onment*	42	-0.277	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	33	8.2	7.8
Quality	Drinking Water Violations (% of community water systems)	48	6.3%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	4	24.1%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	5	24.5%	32.3
Clinical Care*		49	-0.944	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	45	5.0%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	28	30.6	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	27	41.7	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	36	74.7%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	46	62.3%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	48	63.2%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	27	96.1%	96.1
Care	Hospice Care (% of Medicare decedents)	41	42.3%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	49	20.3%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	47	2,062	1,48
Behaviors*		47	-1.312	
Nutrition	Exercise (% of adults ages 65+)	48	13.4%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	44	4.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	42	36.2%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	36	26.8%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	48	14.3%	8.99
Health Outcon	nes*	43	-0.510	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	50	10.7%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	11	7.2	9.9
	Excessive Drinking (% of adults ages 65+)	7	4.8%	7.09
	Frequent Mental Distress (% of adults ages 65+)	18	7.5%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	29	18.4	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	50	3,147	2,15
•	Early Death Racial Disparity (ratio)***	_	1.2	1.5
Physical Health	Falls (% of adults ages 65+)	14	26.2%	27.19
•	Frequent Physical Distress (% of adults ages 65+)	44	19.4%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	44	58%	52%
	Obesity (% of adults ages 65+)	30	31.5%	29.5
	Teeth Extractions (% of adults ages 65+)	47	20.0%	13.49
Overall	, , , , , , , , , , , , , , , , , , , ,	50	-1.000	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of excessive drinking
- High volunteerism rate
- · Low prevalence of avoiding care due to cost

Challenges

- · High early death rate
- · Low percentage of households with high-speed internet
- Low prevalence of fruit and vegetable consumption

Highlights

Number of Geriatric Providers

from 31.4 to 36.5 per 100,000 adults ages 65+ between September 2020 and September 2022.

Poverty

from 8.2% to 9.5% of adults ages 65+ between 2018 and 2021.

Multiple Chronic Conditions

from 47% to 53% of Medicare beneficiaries ages 65+ between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	41	-0.404	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	31	16.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	41	16.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	41	14.0%	11.9%
Resources	Poverty (% of adults ages 65+)	25	9.5%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	32	58.2	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	24	\$42	\$62
and	High-speed Internet (% of households with adults ages 65+)	36	80.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	49	31.3%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	33	60	_
	Volunteerism (% of adults ages 65+)	17	27.6%	22.1%
Physical Enviro	onment*	17	0.437	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	25	7.6	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	17	26.8%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	20	27.3%	32.3%
Clinical Care*		20	0.175	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	6	2.2%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	19	36.5	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	10	65.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	29	76.8%	75.9%
linical	Flu Vaccination (% of adults ages 65+)	19	69.9%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	16	72.9%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	23	96.3%	96.1%
Care	Hospice Care (% of Medicare decedents)	24	47.5%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	43	25.6%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	42	1,796	1,482
Behaviors*		44	-0.834	
Nutrition	Exercise (% of adults ages 65+)	44	15.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	48	3.8%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	37	34.7%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	23	25.2%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	36	10.3%	8.9%
Health Outcon		41	-0.439	
3ehavioral	Cognitive Difficulty (% of adults ages 65+)‡	33	8.3%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	19	8.8	9.9
	Excessive Drinking (% of adults ages 65+)	18	6.6%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	34	8.6%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	27	18.0	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	37	2,429	2,151
	Early Death Racial Disparity (ratio)***	_	1.4	1.5
Physical Health	Falls (% of adults ages 65+)	41	30.6%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	36	16.5%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	28	53%	52%
	Obesity (% of adults ages 65+)	35	32.5%	29.5%
	Teeth Extractions (% of adults ages 65+)	43	17.7%	13.4%
Overall	•	40	-0.341	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of multiple chronic conditions
- Low prevalence of food insecurity
- Low prevalence of avoiding care due to cost

Challenges

- · High suicide rate
- High prevalence of excessive drinking
- Low geriatric provider rate

Highlights

Obesity

28%

from 24.1% to 30.9% of adults ages 65+ between 2018 and 2021.

High-speed Internet

23%

from 67.8% to 83.3% of households with adults ages 65+ between 2016 and 2021.

Home Health Care Workers

23%

from 32.5 to 40.1 aides per 1,000 adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	26	0.174	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	49	27.3	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	35	15.0	11.8
Economic	Food Insecurity (% of adults ages 60+)	9	9.2%	11.9%
Resources	Poverty (% of adults ages 65+)	15	8.8%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	33	57.9	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	12	\$83	\$62
ind	High-speed Internet (% of households with adults ages 65+)	21	83.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	45	24.8%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	12	39	_
	Volunteerism (% of adults ages 65+)	18	27.1%	22.1%
Physical Enviro	onment*	6	0.757	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	8	6.1	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	10	25.4%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	19	27.2%	32.3%
Clinical Care*	,	41	-0.356	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	14	2.6%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	46	21.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	31	40.1	60.3
Clinical	Cancer Screenings (% of adults ages 65-75)	38	74.4%	75.9%
	Flu Vaccination (% of adults ages 65+)	36	65.0%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	36	69.3%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	44	93.4%	96.1%
Quality of Care	Hospice Care (% of Medicare decedents)	44	41.2%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	36	29.4%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	5	929	1,482
Behaviors*	,	10	0.422	
Nutrition	Exercise (% of adults ages 65+)	5	28.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	31	5.9%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or	20	29.6%	31.29
	better health)			
Sleep Health	Insufficient Sleep (% of adults ages 65+)	5	21.2%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	28	9.0%	8.9%
Health Outcon	nes*	28	-0.036	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	11	6.7%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	5	6.0	9.9
	Excessive Drinking (% of adults ages 65+)	42	8.6%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	24	7.8%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	49	34.1	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	23	1,980	2,151
	Early Death Racial Disparity (ratio)***		2.1	1.5
Physical Health	Falls (% of adults ages 65+)	39	29.9%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	25	15.6%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	3	38%	52%
	Obesity (% of adults ages 65+)	25	30.9%	29.5%
	Teeth Extractions (% of adults ages 65+)	23	12.9%	13.49
	-	23	0.094	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.
Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.
Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of frequent mental distress
- · Low prevalence of food insecurity
- · Low prevalence of avoiding care due to cost

Challenges

- High prevalence of obesity
- · Low percentage of households with high-speed internet
- · Low home health care worker rate

Highlights

Poverty

from 7.2% to 8.7% of adults ages 65+ between 2017 and 2021.

Obesity

from 29.9% to 34.9% of adults ages 65+ between 2016 and 2021.

Number of Geriatric Providers

from 23.8 to 26.4 per 100,000 adults ages 65+ between September 2019 and September 2022.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	22	0.225	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	12	11.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	31	14.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	7	8.5%	11.9%
Resources	Poverty (% of adults ages 65+)	13	8.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	39	52.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	15	\$57	\$62
and	High-speed Internet (% of households with adults ages 65+)	34	80.6%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	33	16.9%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	16	43	_
	Volunteerism (% of adults ages 65+)	13	29.8%	22.1%
Physical Enviro	onment*	5	0.777	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	10	6.2	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	24	28.1%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	15	26.8%	32.3%
Clinical Care*		13	0.405	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	1	1.7%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	36	26.4	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	34	34.2	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	42	73.7%	75.9%
linical ervices	Flu Vaccination (% of adults ages 65+)	16	71.1%	67.5%
services	Pneumonia Vaccination (% of adults ages 65+)	5	75.3%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	21	96.4%	96.1%
Care	Hospice Care (% of Medicare decedents)	18	48.3%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	8	43.7%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	14	1,161	1,482
Behaviors*		17	0.200	
Nutrition	Exercise (% of adults ages 65+)	26	22.7%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	22	7.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	34	33.4%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	5	21.2%	26.0%
Гobacco Use	Smoking (% of adults ages 65+)	16	8.2%	8.9%
Health Outcon	nes*	12	0.390	
3ehavioral	Cognitive Difficulty (% of adults ages 65+)‡	11	6.7%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	1	3.9	9.9
	Excessive Drinking (% of adults ages 65+)	24	7.0%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	4	6.3%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	10	13.9	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	19	1,945	2,151
	Early Death Racial Disparity (ratio)***		2.1	1.5
Physical Health	Falls (% of adults ages 65+)	19	27.6%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	6	12.4%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	19	48%	52%
	Obesity (% of adults ages 65+)	44	34.9%	29.5%
	Teeth Extractions (% of adults ages 65+)	9	10.6%	13.4%
Overall		13	0.325	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of obesity
- · Low prevalence of excessive drinking
- · Low prevalence of physical inactivity

Challenges

- · High suicide rate
- · High poverty rate
- · Low flu vaccination rate

Highlights

Home Health Care Workers

from 30.7 to 26.0 aides per 1,000 adults ages 65+ between 2018 and 2021.

High-speed Internet

from 74.8% to 85.6% of households with adults ages 65+ between 2016 and 2021.

Early Deaths

from 2,234 to 2,403 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

Measure	es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	34	-0.118	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)+	48	25.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	17	10.6	11.8
Economic	Food Insecurity (% of adults ages 60+)	23	10.8%	11.9%
Resources	Poverty (% of adults ages 65+)	41	11.3%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	15	91.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	40	\$30	\$62
and .	High-speed Internet (% of households with adults ages 65+)	12	85.6%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	28	15.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	30	56	_
	Volunteerism (% of adults ages 65+)	43	18.0%	22.1%
Physical Enviro	onment*	47	-0.487	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	47	9.6	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	44	35.2%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	42	35.9%	32.3%
Clinical Care*		48	-0.790	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	43	4.7%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	24	33.6	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	44	26.0	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	24	77.7%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	48	60.2%	67.5%
Sei vices	Pneumonia Vaccination (% of adults ages 65+)	46	64.3%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	46	91.9%	96.1%
Care	Hospice Care (% of Medicare decedents)	29	46.4%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	33	30.7%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	30	1,514	1,482
Behaviors*		42	-0.734	
Nutrition	Exercise (% of adults ages 65+)	33	20.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	39	4.7%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	16	28.3%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	43	29.4%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	46	12.6%	8.9%
Health Outcon	nes*	40	-0.403	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	36	8.5%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	47	16.4	9.9
	Excessive Drinking (% of adults ages 65+)	12	6.0%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	46	11.5%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	48	30.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	36	2,403	2,151
	Early Death Racial Disparity (ratio)***		1.5	1.5
Physical Health	Falls (% of adults ages 65+)	25	28.3%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	46	20.5%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	16	47%	52%
	Obesity (% of adults ages 65+)	6	25.1%	29.5%
	Teeth Extractions (% of adults ages 65+)	26	13.0%	13.4%
Overall		43	-0.465	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

New Hampshire

Overall Rank

Summary

Strengths

- Low prevalence of frequent physical distress
- · Low prevalence of food insecurity
- · Low prevalence of avoiding care due to cost

Challenges

- · High prevalence of excessive drinking
- · Low home health care worker rate
- · High prevalence of severe housing problems

Highlights

Poverty

from 4.6% to 6.9% of adults ages 65+ between 2016 and 2021.

Number of Geriatric Providers

from 34.4 to 40.7 per 100,000 adults ages 65+ between September 2019 and September 2022.

High-speed Internet

from 82.1% to 87.4% of households with adults ages 65+ between 2017 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	4	0.768	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	14	12.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	13	10.3	11.8
Economic	Food Insecurity (% of adults ages 60+)	1	6.4%	11.99
Resources	Poverty (% of adults ages 65+)	2	6.9%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	47	42.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	14	\$66	\$62
and	High-speed Internet (% of households with adults ages 65+)	5	87.4%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	40	18.4%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	2	19	_
	Volunteerism (% of adults ages 65+)	13	29.8%	22.19
Physical Enviro	onment*	9	0.540	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	2	4.6	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)*	38	32.4%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	42	35.9%	32.3
Clinical Care*		6	0.647	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	1	1.7%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	13	40.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	40	29.9	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	4	81.3%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	3	74.9%	67.5
Dei vices	Pneumonia Vaccination (% of adults ages 65+)	8	75.1%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	9	97.0%	96.1
Care	Hospice Care (% of Medicare decedents)	18	48.3%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	31	31.6%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	18	1,269	1,48
Behaviors*		4	0.968	
Nutrition	Exercise (% of adults ages 65+)	11	27.2%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	4	11.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	8	26.4%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	19	23.9%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	7	7.4%	8.99
Health Outcon	nes*	5	0.558	
3ehavioral	Cognitive Difficulty (% of adults ages 65+)*	4	6.0%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	3	5.6	9.9
	Excessive Drinking (% of adults ages 65+)	34	7.8%	7.09
	Frequent Mental Distress (% of adults ages 65+)	6	7.0%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	29	18.4	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	8	1,743	2,15
SI 1 111 121	Early Death Racial Disparity (ratio)***	_	1.2	1.5
Physical Health	Falls (% of adults ages 65+)	28	28.5%	27.19
	Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	5 11	12.1% 44%	15.8° 52%
	ages 65-74)			
	Obesity (% of adults ages 65+)	7	25.9%	29.5
	Teeth Extractions (% of adults ages 65+)	12	11.0%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

New Jersey State Health Department Website: nj.gov/health/

Overall Rank

Summary

Strengths

- · Low early death rate
- · Low prevalence of obesity
- · High geriatric provider rate

Challenges

- · High prevalence of multiple chronic conditions
- · Low hospice care use
- · High prevalence of severe housing problems

Highlights

Drug Deaths

from 4.0 to 10.3 deaths per 100,000 adults ages 65+ between 2014-2016 and 2019-2021.

Flu Vaccination Rates

from 57.2% to 67.1% of adults ages 65+ between 2013 and 2021.

High-speed Internet

from 73.6% to 85.1% of households with adults ages 65+ between 2016 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	25	0.181	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	3	4.4	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	7	7.8	11.8
Conomic	Food Insecurity (% of adults ages 60+)	17	9.7%	11.9%
Resources	Poverty (% of adults ages 65+)	22	9.4%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	14	94.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	24	\$42	\$62
nd	High-speed Internet (% of households with adults ages 65+)	15	85.1%	83.1%
ngagement	Low-care Nursing Home Residents (% of residents)	27	15.6%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	28	52	_
	Volunteerism (% of adults ages 65+)	38	19.1%	22.1%
Physical Enviro	onment*	50	-1.333	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	28	7.8	7.8
Quality	Drinking Water Violations (% of community water systems)	49	8.9%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	49	40.0%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	50	43.6%	32.3%
Clinical Care*		38	-0.280	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	41	4.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	5	49.2	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	20	46.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	40	74.0%	75.9%
linical	Flu Vaccination (% of adults ages 65+)	26	67.1%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	47	64.2%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	32	95.7%	96.1%
Care	Hospice Care (% of Medicare decedents)	46	39.5%	46.79
Care	Nursing Home Quality (% of beds rated four or five stars)	11	41.5%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	28	1,472	1,482
Behaviors*		26	0.036	
Nutrition	Exercise (% of adults ages 65+)	_	_	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	2	12.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	29	32.4%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	42	28.6%	26.09
obacco Use	Smoking (% of adults ages 65+)	16	8.2%	8.9%
Health Outcon	nes*	15	0.323	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	11	6.7%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	28	10.3	9.9
	Excessive Drinking (% of adults ages 65+)	21	6.8%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	39	9.4%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	2	9.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	6	1,727	2,151
	Early Death Racial Disparity (ratio)***	_	1.6	1.5
hysical Health	Falls (% of adults ages 65+)	6	25.0%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	31	16.3%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	40	56%	52%
	Obesity (% of adults ages 65+)	8	26.3%	29.5%
	Teeth Extractions (% of adults ages 65+)	9	10.6%	13.4%
Overall		29	0.020	1

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- · Low prevalence of multiple chronic conditions
- · Low prevalence of excessive drinking
- · High home health care worker rate

Challenges

- · High suicide rate
- · High poverty rate
- · Low percentage of older adults with a dedicated health care provider

Highlights

Drug Deaths

from 7.2 to 13.7 deaths per 100,000 adults ages 65+ between 2015-2017 and 2019-2021.

Early Deaths

from 1,694 to 2,183 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

High-speed Internet

from 63.7% to 78.2% of households with adults ages 65+ between 2016 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	47	-0.890	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	46	22.7	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	32	14.6	11.8
Economic	Food Insecurity (% of adults ages 60+)	36	13.5%	11.99
Resources	Poverty (% of adults ages 65+)	48	12.8%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	31	60.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	20	\$50	\$62
and .	High-speed Internet (% of households with adults ages 65+)	45	78.2%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	44	22.2%	15.2
	Risk of Social Isolation (index from 1-100, adults ages 65+)	48	91	_
	Volunteerism (% of adults ages 65+)	22	25.3%	22.19
Physical Enviro	onment*	25	0.230	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	13	6.7	7.8
Quality	Drinking Water Violations (% of community water systems)	45	2.9%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	19	27.1%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	17	27.0%	32.3
Clinical Care*		40	-0.338	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	39	4.0%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	37	26.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	5	85.9	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	45	71.7%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	28	66.8%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	39	68.6%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	47	91.4%	96.1
Care	Hospice Care (% of Medicare decedents)	34	45.4%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	39	28.6%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	9	1,032	1,48
Behaviors*		36	-0.198	
Nutrition	Exercise (% of adults ages 65+)	16	25.2%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	44	4.2%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	22	30.0%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	34	26.7%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	32	9.2%	8.99
Health Outcon	nes*	26	-0.019	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	45	9.5%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	44	13.7	9.9
	Excessive Drinking (% of adults ages 65+)	4	4.4%	7.09
	Frequent Mental Distress (% of adults ages 65+)	39	9.4%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	47	26.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	30	2,183	2,15
	Early Death Racial Disparity (ratio)***	_	1.6	1.5
Physical Health	Falls (% of adults ages 65+)	48	33.2%	27.19
,	Frequent Physical Distress (% of adults ages 65+)	31	16.3%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	9	43%	529
	Obesity (% of adults ages 65+)	11	26.7%	29.5
	Teeth Extractions (% of adults ages 65+)	23	12.9%	13.4
Overall		41	-0.367	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- · Low suicide rate
- · High level of community support expenditures
- · High home health care worker rate

Challenges

- · High poverty rate
- High risk of social isolation
- · Low hospice care use

Highlights

Drug Deaths

from 6.7 to 9.6 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

High-speed Internet

from 70.4% to 82.6% of households with adults ages 65+ between 2016 and 2021.

Early Deaths

from 2,026 to 1,737 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	35	-0.158	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	4	4.5	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	2	6.4	11.8
conomic	Food Insecurity (% of adults ages 60+)	30	12.4%	11.99
Resources	Poverty (% of adults ages 65+)	47	12.2%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.C
Social Support	Community Support Expenditures (dollars per adult ages 60+)	9	\$100	\$62
ind	High-speed Internet (% of households with adults ages 65+)	27	82.6%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	6	9.5%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	47	86	_
	Volunteerism (% of adults ages 65+)	36	19.8%	22.19
Physical Enviro	onment*	40	-0.153	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	11	6.5	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	48	39.7%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	48	41.1%	32.3
Clinical Care*		29	-0.021	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	35	3.5%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	16	38.5	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	1	137.6	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	14	79.0%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	28	66.8%	67.5
bei vices	Pneumonia Vaccination (% of adults ages 65+)	43	65.6%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	30	96.0%	96.1
Care	Hospice Care (% of Medicare decedents)	50	24.7%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	25	34.9%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	27	1,468	1,48
Behaviors*		13	0.336	
Nutrition	Exercise (% of adults ages 65+)	20	24.5%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	3	12.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	26	31.2%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	43	29.4%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	16	8.2%	8.99
Health Outcon	nes*	8	0.431	
3ehavioral	Cognitive Difficulty (% of adults ages 65+)*	31	7.8%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	21	9.6	9.9
	Excessive Drinking (% of adults ages 65+)	23	6.9%	7.09
	Frequent Mental Distress (% of adults ages 65+)	29	8.3%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	3	9.6	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	7	1,737	2,15
	Early Death Racial Disparity (ratio)***		1.4	1.5
Physical Health	Falls (% of adults ages 65+)	8	25.1%	27.19
	Frequent Physical Distress (% of adults ages 65+)	18	14.5%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	30	54%	52%
	Obesity (% of adults ages 65+)	14	27.1%	29.5
	Teeth Extractions (% of adults ages 65+)	19	12.0%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

North Carolina

State Health Department Website: ncdhhs.gov

Summary

Strengths

- · High level of community support expenditures
- · High pneumonia vaccination rate
- · Low prevalence of physical inactivity

Challenges

- · High prevalence of obesity
- Low percentage of households with high-speed internet
- · Low home health care worker rate

Highlights

Drug Deaths

from 3.8 to 7.2 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Food Insecurity

from 17.2% to 13.7% of adults ages 60+ between 2017-2018 and 2019-2020.

Poverty

from 8.7% to 10.2% of adults ages 65+ between 2018 and 2021.

rice Factors* Firearm Deaths (deaths per 100,000 adults ages 65+)* Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	28	0.123	
Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)+	20	14.0	13.1
	25	12.9	11.8
Food Insecurity (% of adults ages 60+)	38	13.7%	11.9%
Poverty (% of adults ages 65+)	33	10.2%	10.39
SNAP Reach (participants per 100 adults ages 60+ in poverty)	24	75.9	81.0
Community Support Expenditures (dollars per adult ages 60+)	5	\$138	\$62
High-speed Internet (% of households with adults ages 65+)	34	80.6%	83.19
Low-care Nursing Home Residents (% of residents)	11	10.7%	15.29
Risk of Social Isolation (index from 1-100, adults ages 65+)	29	55	_
Volunteerism (% of adults ages 65+)	32	22.2%	22.19
onment*	16	0.440	
Air Pollution (micrograms of fine particles per cubic meter)	18	7.2	7.8
	1	0.0%	0.89
Housing Cost Burden (% of households with adults	27	28.7%	31.89
Severe Housing Problems (% of small households with	25	29.1%	32.3
,	21	0.162	
Avoided Care Due to Cost (% of adults ages 65+)	36	3.6%	3.49
Geriatric Providers (providers per 100,000 adults ages 65+)	18	37.0	36.4
Home Health Care Workers (workers per 1,000 adults ages 65+)	38	31.6	60.3
Cancer Screenings (% of adults ages 65-75)	5	80.4%	75.9
Flu Vaccination (% of adults ages 65+)	19	69.9%	67.5
Pneumonia Vaccination (% of adults ages 65+)	7	75.2%	69.7
Dedicated Health Care Provider (% of adults ages 65+)	7	97.1%	96.19
Hospice Care (% of Medicare decedents)	23	47.9%	46.7
Nursing Home Quality (% of beds rated four or five stars)	42	26.9%	32.6
Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	35	1,609	1,48
	37	-0.212	
Exercise (% of adults ages 65+)	42	19.1%	23.19
Fruit and Vegetable Consumption (% of adults ages 65+)	43	4.3%	7.3%
Physical Inactivity (% of adults ages 65+ in fair or better health)	12	27.5%	31.29
Insufficient Sleep (% of adults ages 65+)	24	25.3%	26.0
Smoking (% of adults ages 65+)	23	8.5%	8.9%
nes*	37	-0.263	
Cognitive Difficulty (% of adults ages 65+)*	27	7.6%	7.89
Drug Deaths (deaths per 100,000 adults ages 65+)*	11	7.2	9.9
Excessive Drinking (% of adults ages 65+)	24	7.0%	7.09
Frequent Mental Distress (% of adults ages 65+)	25	7.9%	8.59
Suicide (deaths per 100,000 adults ages 65+)	17	16.1	16.9
Early Death (deaths per 100,000 adults ages 65-74)	35	2,310	2,15
Early Death Racial Disparity (ratio)***		1.3	1.5
Falls (% of adults ages 65+)	27	28.4%	27.19
Frequent Physical Distress (% of adults ages 65+)	37	16.6%	15.89
Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	30	54%	52%
Obesity (% of adults ages 65+)	38	33.0%	29.5
Teeth Extractions (% of adults ages 65+)	35	15.8%	13.49
	Community Support Expenditures (dollars per adult ages 60+) High-speed Internet (% of households with adults ages 65+) Low-care Nursing Home Residents (% of residents) Risk of Social Isolation (index from 1-100, adults ages 65+) Volunteerism (% of adults ages 65+) ### Air Pollution (micrograms of fine particles per cubic meter) Drinking Water Violations (% of community water systems) Housing Cost Burden (% of households with adults ages 65+) Severe Housing Problems (% of small households with adults ages 65+) Geriatric Providers (providers per 100,000 adults ages 65+) Home Health Care Workers (workers per 1,000 adults ages 65+) Cancer Screenings (% of adults ages 65-75) Flu Vaccination (% of adults ages 65+) Dedicated Health Care Provider (% of adults ages 65+) Hospice Care (% of Medicare decedents) Nursing Home Quality (% of beds rated four or five stars) Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74) Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+) Smoking (% of adults ages 65+) Fruit peaths (deaths per 100,000 adults ages 65+) ##### Brug Deaths (deaths per 100,000 adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Early Death (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74) Obesity (% of adults ages 65+)	Community Support Expenditures (dollars per adult ages 60+) High-speed Internet (% of households with adults ages 65+) 134 Low-care Nursing Home Residents (% of residents) 118 Risk of Social Isolation (index from 1-100, adults ages 65+) 29 Volunteerism (% of adults ages 65+) 316 Air Pollution (micrograms of fine particles per cubic meter) 18 Drinking Water Violations (% of community water systems) 19 Housing Cost Burden (% of households with adults ages 65+) 40 Severe Housing Problems (% of small households with adults ages 65+) 41 Severe Housing Problems (% of adults ages 65+) 42 Avoided Care Due to Cost (% of adults ages 65+) 43 Geriatric Providers (providers per 100,000 adults ages 65+) Home Health Care Workers (workers per 1,000 adults ages 65+) Cancer Screenings (% of adults ages 65-75) 51 Flu Vaccination (% of adults ages 65+) 72 Dedicated Health Care Provider (% of adults ages 65+) Pneumonia Vaccination (% of adults ages 65+) 73 Dedicated Health Care Provider (% of adults ages 65+) Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74) Exercise (% of adults ages 65-74) Exercise (% of adults ages 65+) Fruit and Vegetable Consumption (% of adults ages 65+) Physical Inactivity (% of adults ages 65+) Smoking (% of adults ages 65+) Cognitive Difficulty (% of adults ages 65+) Exercise (% of adults ages 65+) Exercise (% of adults ages 65+) Cognitive Difficulty (% of adults ages 65+) Exercise (% of adults ages 65+) Fruquent Mental Distress (% of adults ages 65+) Excessive Drinking (% of adults ages 65+) Ex	Community Support Expenditures (dollars per adult ages 60+) 5 \$138 High-speed Internet (% of households with adults ages 65+) 34 80.6% Low-care Nursing Home Residents (% of residents) 11 10.7% Risk of Social Isolation (index from 1-100, adults ages 65+) 29 55 Volunteerism (% of adults ages 65+) 32 22.2% Imment* 16 0.440 Air Pollution (micrograms of fine particles per cubic meter) 18 7.2 Drinking Water Violations (% of community water systems) 1 0.0% Housing Coost Burden (% of households with adults ages 65+)* 28.7% 28.7% Severe Housing Problems (% of small households with adults ages 65+)* 25 29.1% Severe Housing Problems (% of adults ages 65+) 36 3.6% Geriatric Providers (providers per 100,000 adults ages 65+) 18 37.0 Home Health Care Workers (workers per 1,000 adults ages 65+) 18 37.0 Home Health Care Workers (workers per 1,000 adults ages 65+) 19 69.9% Flu Vaccination (% of adults ages 65+) 7 75.2% Dedicated Health Care Provider (% of adults ages 65+)

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

North Dakota

Overall Rank

Summary

Strengths

- Low prevalence of frequent mental distress
- Low prevalence of food insecurity
- · Low prevalence of avoiding care due to cost

Challenges

- · High prevalence of obesity
- · High prevalence of excessive drinking
- · Low hospice care use

Highlights

Number of Geriatric Providers

from 13.0 to 21.7 per 100,000 adults ages 65+ between September 2021 and September 2022.

High-speed Internet

from 64.6% to 79.1% of households with adults ages 65+ between 2016 and 2021.

Home Health Care Workers

from 58.0 to 50.1 aides per 1,000 adults ages 65+ between 2016 and 2021.

Measure	es es	State Rank	State Value	U.S Valu
Social & Econo	omic Factors*	16	0.330	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	14	12.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	37	15.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	2	6.5%	11.99
Resources	Poverty (% of adults ages 65+)	22	9.4%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	46	44.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	8	\$102	\$62
and	High-speed Internet (% of households with adults ages 65+)	43	79.1%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	25	15.4%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	17	45	_
	Volunteerism (% of adults ages 65+)	7	33.2%	22.19
Physical Enviro	onment*	2	1.253	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	4	5.0	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	5	24.6%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	3	23.3%	32.3
Clinical Care*		31	-0.067	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	3	1.9%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	43	21.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	14	50.1	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	32	76.2%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	23	69.5%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	30	70.4%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	41	94.2%	96.1
Care	Hospice Care (% of Medicare decedents)	48	31.6%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	3	48.9%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	26	1,452	1,48
Behaviors*		18	0.186	
Nutrition	Exercise (% of adults ages 65+)	37	20.3%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	22	7.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	24	30.8%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	9	22.6%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	8	7.5%	8.99
Health Outcon	nes*	11	0.397	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	2	5.7%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	_	_	9.9
	Excessive Drinking (% of adults ages 65+)	37	8.0%	7.09
	Frequent Mental Distress (% of adults ages 65+)	1	5.7%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	7	13.1	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	17	1,907	2,15
	Early Death Racial Disparity (ratio)***	_	2.1	1.5
Physical Health	Falls (% of adults ages 65+)	12	26.0%	27.19
	Frequent Physical Distress (% of adults ages 65+)	6	12.4%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	19	48%	52%
	Obesity (% of adults ages 65+)	47	35.6%	29.5
	Teeth Extractions (% of adults ages 65+)	17	11.7%	13.4

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Summary

Strengths

- · Low prevalence of excessive drinking
- · Low poverty rate
- · High hospice care use

Challenges

- · High early death rate
- High prevalence of obesity
- · High prevalence of physical inactivity

Highlights

Drug Deaths

from 7.3 to 10.5 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Number of Geriatric Providers

from 32.9 to 41.0 per 100,000 adults ages 65+ between September 2019 and September 2022.

High-speed Internet

from 68.6% to 80.2% of households with adults ages 65+ between 2016 and 2021.

Measure	es e	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	32	-0.046	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	18	13.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	15	10.4	11.8
Economic	Food Insecurity (% of adults ages 60+)	22	10.2%	11.9%
Resources	Poverty (% of adults ages 65+)	25	9.5%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	21	78.6	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	31	\$36	\$62
and	High-speed Internet (% of households with adults ages 65+)	37	80.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	29	15.9%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	37	63	_
	Volunteerism (% of adults ages 65+)	23	24.9%	22.1%
Physical Enviro		29	0.153	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	41	8.7	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults	21	27.6%	31.8%
-	ages 65+)‡			
	Severe Housing Problems (% of small households with adults ages 62+)	23	28.4%	32.3%
Clinical Care*		22	0.158	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	22	2.8%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	11	41.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	22	45.3	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	28	77.0%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	32	66.3%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	27	70.7%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	9	97.0%	96.1%
Care	Hospice Care (% of Medicare decedents)	3	54.8%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	34	30.6%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	37	1,713	1,482
Behaviors*	,	40	-0.684	
Nutrition	Exercise (% of adults ages 65+)	35	20.5%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	20	5.0%	7.3%
		30		
	Physical Inactivity (% of adults ages 65+ in fair or better health)	38 40	35.6%	31.2%
Activity	Physical Inactivity (% of adults ages 65+ in fair or			
Activity Sleep Health	Physical Inactivity (% of adults ages 65+ in fair or better health)	40	35.6%	31.2% 26.0% 8.9%
Sleep Health Tobacco Use	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	40 30	35.6% 26.1%	26.0%
Activity Sleep Health Tobacco Use Health Outcon	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes*	30 37	35.6% 26.1% 11.0%	26.0%
Activity Sleep Health Tobacco Use Health Outcon Behavioral	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+)	30 37 39	35.6% 26.1% 11.0% - 0.304	26.0% 8.9%
Activity Sleep Health Tobacco Use Health Outcon Behavioral	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+)	30 37 39 22 32	35.6% 26.1% 11.0% - 0.304 7.4% 10.5	26.0% 8.9% 7.8% 9.9
Activity Sleep Health Tobacco Use Health Outcon Behavioral	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+)	30 37 39 22 32 9	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2%	26.0% 8.9% 7.8% 9.9 7.0%
Activity Sleep Health Tobacco Use Health Outcon Behavioral	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	30 37 39 22 32	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9%	26.0% 8.9% 7.8% 9.9 7.0% 8.5%
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	30 37 39 22 32 9 38 17	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1	7.8% 9.9 7.0% 8.5% 16.9
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	30 37 39 22 32 9 38	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488	7.8% 9.9 7.0% 8.5% 16.9 2,151
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***	30 37 39 22 32 9 38 17 39	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488 1.4	7.8% 9.9 7.0% 8.5% 16.9 2,151 1.5
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) nes* Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+)	30 37 39 22 32 9 38 17 39 -	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488 1.4 28.0%	26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality Physical Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) mes* Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	30 37 39 22 32 9 38 17 39	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488 1.4	7.8% 9.9 7.0% 8.5% 16.9 2,151
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) Cognitive Difficulty (% of adults ages 65+) Drug Deaths (deaths per 100,000 adults ages 65+) Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	30 37 39 22 32 9 38 17 39 — 22 27 34	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488 1.4 28.0% 15.7% 55%	26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5 27.1% 15.8% 52%
Activity Sleep Health Tobacco Use Health Outcon Behavioral Health Mortality	Physical Inactivity (% of adults ages 65+ in fair or better health) Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+) mes* Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	30 37 39 22 32 9 38 17 39 —	35.6% 26.1% 11.0% -0.304 7.4% 10.5 5.2% 8.9% 16.1 2.488 1.4 28.0% 15.7%	26.0% 8.9% 7.8% 9.9 7.0% 8.5% 16.9 2.151 1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Strengths

- · Low prevalence of excessive drinking
- · High pneumonia vaccination rate
- · Low prevalence of severe housing problems

Challenges

- · High prevalence of food insecurity
- · Low home health care worker rate
- · High prevalence of physical inactivity

Highlights

Number of Geriatric Providers

from 15.7 to 21.7 per 100,000 adults ages 65+ between September 2019 and September 2022.

Early Deaths

from 2,379 to 2,975 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Poverty

from 8.7% to 10.6% of adults ages 65+ between 2018 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	46	-0.812	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	45	21.0	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	34	14.9	11.8
Economic	Food Insecurity (% of adults ages 60+)	47	17.7%	11.9%
Resources	Poverty (% of adults ages 65+)	35	10.6%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	28	63.6	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	44	\$26	\$62
and	High-speed Internet (% of households with adults ages 65+)	39	79.9%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	48	28.3%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	42	76	_
	Volunteerism (% of adults ages 65+)	24	24.8%	22.1%
Physical Enviro	onment*	34	0.033	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	41	8.7	7.8
Quality	Drinking Water Violations (% of community water systems)	44	2.4%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	12	25.6%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	7	24.6%	32.39
Clinical Care*		42	-0.428	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	40	4.3%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	43	21.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	43	26.2	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	46	71.5%	75.99
Clinical Services	Flu Vaccination (% of adults ages 65+)	25	68.0%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	5	75.3%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	32	95.7%	96.19
Care	Hospice Care (% of Medicare decedents)	18	48.3%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	40	27.9%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	44	1,829	1,482
Behaviors*		45	-1.178	
Nutrition	Exercise (% of adults ages 65+)	46	14.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	49	3.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	45	39.0%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	12	23.1%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	43	12.2%	8.9%
Health Outcon	nes*	45	-0.677	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	43	9.4%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	26	10.1	9.9
	Excessive Drinking (% of adults ages 65+)	3	4.3%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	41	9.5%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	41	23.3	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	48	2,975	2,151
	Early Death Racial Disparity (ratio)***	_	1.3	1.5
Physical Health	Falls (% of adults ages 65+)	33	29.4%	27.1%
,	Frequent Physical Distress (% of adults ages 65+)	37	16.6%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	40	56%	52%
	Obesity (% of adults ages 65+)	40	33.2%	29.59
	Teeth Extractions (% of adults ages 65+)	45	19.2%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

State Health Department Website: oregon.gov/oha/ph

Overall Rank

Summary

Strengths

- · Low prevalence of multiple chronic conditions
- · High pneumonia vaccination rate
- · Low prevalence of physical inactivity

Challenges

- · High suicide rate
- · High prevalence of excessive drinking
- Low percentage of older adults with a dedicated health care provider

Highlights

Drug Deaths

from 5.9 to 11.2 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Fruit and Vegetable Consumption

from 12.5% to 5.5% of adults ages 65+ between 2017 and 2021.

High-speed Internet

from 76.0% to 86.1% of households with adults ages 65+ between 2016 and 2021.

Measure	es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	10	0.479	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	37	19.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	27	13.3	11.8
Economic	Food Insecurity (% of adults ages 60+)	15	9.6%	11.9%
Resources	Poverty (% of adults ages 65+)	20	9.3%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	30	\$37	\$62
ind	High-speed Internet (% of households with adults ages 65+)	9	86.1%	83.1%
ingagement	Low-care Nursing Home Residents (% of residents)	7	9.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	19	46	_
	Volunteerism (% of adults ages 65+)	27	23.8%	22.1%
Physical Enviro		44	-0.360	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	44	8.9	7.8
Quality	Drinking Water Violations (% of community water systems)	30	0.3%	0.8%
Housing	Housing Cost Burden (% of households with adults	43	34.4%	31.8%
.	ages 65+)*			
	Severe Housing Problems (% of small households with adults ages 62+)	41	35.7%	32.3%
Clinical Care*		23	0.154	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	24	2.9%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	35	27.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	23	44.9	60.3
reventive	Cancer Screenings (% of adults ages 65-75)	25	77.5%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	33	66.0%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	2	75.5%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	45	92.9%	96.1%
	Hospice Care (% of Medicare decedents)	6	54.5%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	25	34.9%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	7	998	1,482
Behaviors*	•	9	0.554	
Nutrition	Exercise (% of adults ages 65+)	14	26.3%	23.1%
nd Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	36	5.5%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	2	22.0%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	17	23.4%	26.0%
obacco Use	Smoking (% of adults ages 65+)	16	8.2%	8.9%
lealth Outcon	nes*	19	0.237	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	36	8.5%	7.8%
lealth	Drug Deaths (deaths per 100,000 adults ages 65+)*	37	11.2	9.9
	Excessive Drinking (% of adults ages 65+)	43	8.8%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	19	7.6%	8.5%
		46	26.1	16.9
Mortality	Suicide (deaths per 100,000 adults ages 65+)	46	26.1 1,968	
Nortality	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)			2,151
	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***	46 21 —	1,968 1.7	2,151 1.5
	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+)	46 21 — 24	1,968 1.7 28.2%	2,151 1.5 27.1%
	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	46 21 —	1,968 1.7	2,151 1.5 27.1%
	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	46 21 — 24 19 7	1,968 1.7 28.2% 14.6% 42%	2,151 1.5 27.1% 15.8% 52%
Mortality Physical Health	Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	46 21 — 24 19	1,968 1.7 28.2% 14.6%	2,151 1.5 27.1% 15.8%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Pennsylvania

Overall Rank

State Health Department Website: health.pa.gov

Summary

Strengths

- · High level of community support expenditures
- High percentage of older adults with a dedicated health care provider
- High geriatric provider rate

Challenges

- High prevalence of multiple chronic conditions
- · High prevalence of frequent physical distress
- · Low percentage of households with high-speed internet

Highlights

Drug Deaths

from 4.3 to 10.3 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Food Insecurity

from 16.5% to 10.9% of adults ages 60+ between 2015-2016 and 2019-2020.

Early Deaths

from 1,775 to 2,153 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	24	0.203	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	13	11.9	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	10	9.1	11.8
Economic	Food Insecurity (% of adults ages 60+)	24	10.9%	11.9%
Resources	Poverty (% of adults ages 65+)	28	9.6%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	11	\$84	\$62
and .	High-speed Internet (% of households with adults ages 65+)	38	80.1%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	21	13.2%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	34	61	_
	Volunteerism (% of adults ages 65+)	20	25.6%	22.19
Physical Enviro	onment*	37	-0.090	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	41	8.7	7.8
Quality	Drinking Water Violations (% of community water systems)	34	0.5%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)†	30	29.5%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	33	31.4%	32.3
Clinical Care*		14	0.402	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	18	2.7%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	9	45.6	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	7	78.5	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	23	78.0%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	16	71.1%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	31	70.3%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	4	97.4%	96.1
Care	Hospice Care (% of Medicare decedents)	31	46.1%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	30	32.2%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	32	1,584	1,48
Behaviors*		35	-0.186	
Nutrition	Exercise (% of adults ages 65+)	29	21.7%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	25	6.9%	7.39
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	28	32.2%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	31	26.3%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	34	9.3%	8.9%
Health Outcon	nes*	36	-0.251	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	15	6.9%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	28	10.3	9.9
	Excessive Drinking (% of adults ages 65+)	32	7.6%	7.09
	Frequent Mental Distress (% of adults ages 65+)	33	8.5%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	15	15.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	28	2,153	2,15
,	Early Death Racial Disparity (ratio)***	_	1.6	1.5
Physical Health	Falls (% of adults ages 65+)	19	27.6%	27.19
,	Frequent Physical Distress (% of adults ages 65+)	41	17.3%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries			
	ages 65-74)	40	56%	52%
	Obesity (% of adults ages 65+)	27	31.0%	29.5
	Teeth Extractions (% of adults ages 65+)	32	14.6%	13.49
Overall		27	0.037	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Rhode Island

Overall Rank

State Health Department Website: health.ri.gov

Summary

Strengths

- · Low suicide rate
- · Low prevalence of obesity
- · High geriatric provider rate

Challenges

- · High prevalence of excessive drinking
- · High risk of social isolation
- · High prevalence of food insecurity

Highlights

Drug Deaths

from 7.7 to 13.3 deaths per 100,000 adults ages 65+ between 2015-2017 and 2019-2021.

High-speed Internet

from 68.2% to 83.2% of households with adults ages 65+ between 2016 and 2021.

Number of Geriatric Providers

from 56.1 to 63.9 per 100,000 adults ages 65+ between September 2019 and September 2022.

<i>A</i> easure	es es	State Rank	State Value	U.S Valu
Social & Econo	omic Factors*	37	-0.232	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	2	4.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	11	9.9	11.8
conomic	Food Insecurity (% of adults ages 60+)	40	13.8%	11.99
Resources	Poverty (% of adults ages 65+)	30	9.8%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	43	\$27	\$62
ind	High-speed Internet (% of households with adults ages 65+)	23	83.2%	83.19
Ingagement	Low-care Nursing Home Residents (% of residents)	30	16.2%	15.2
	Risk of Social Isolation (index from 1-100, adults ages 65+)	40	69	_
	Volunteerism (% of adults ages 65+)	48	16.1%	22.19
Physical Enviro	onment*	36	-0.080	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	16	7.1	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+) [‡]	45	35.4%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	45	38.0%	32.3
Clinical Care*		2	0.936	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	7	2.3%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	1	63.9	36.
	Home Health Care Workers (workers per 1,000 adults ages 65+)	26	41.8	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	2	82.8%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	1	76.5%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	10	74.6%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	1	98.5%	96.1
Care	Hospice Care (% of Medicare decedents)	13	51.2%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	35	29.7%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	20	1,352	1,48
Behaviors*		15	0.214	
Nutrition	Exercise (% of adults ages 65+)	19	24.7%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	11	8.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	18	29.0%	31.2
Sleep Health	Insufficient Sleep (% of adults ages 65+)	45	29.6%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	4	7.0%	8.99
Health Outcon	nes*	20	0.215	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	19	7.1%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)‡	43	13.3	9.9
	Excessive Drinking (% of adults ages 65+)	49	10.6%	7.09
	Frequent Mental Distress (% of adults ages 65+)	28	8.2%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	1	9.0	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	11	1,788	2,15
	Early Death Racial Disparity (ratio)***		1.1	1.5
Physical Health	Falls (% of adults ages 65+)	6	25.0%	27.19
	Frequent Physical Distress (% of adults ages 65+)	29	15.9%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	34	55%	52%
	Obesity (% of adults ages 65+)	4	24.9%	29.5

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

South Carolina

State Health Department Website: scdhec.gov

Summary

Strengths

- · High percentage of older adults with a dedicated health care provider
- · High hospice care use
- · Low prevalence of severe housing problems

Challenges

- · High prevalence of frequent mental distress
- · Low level of community support expenditures
- · Low home health care worker rate

Highlights

Early Deaths

from 1,916 to 2,489 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

High-speed Internet

from 68.0% to 81.3% of households with adults ages 65+ between 2016 and 2021.

Obesity

from 27.1% to 32.5% of adults ages 65+ between 2014 and 2021.

Measure	2 S	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	38	-0.255	
	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	33	16.1	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	43	17.0	11.8
Economic	Food Insecurity (% of adults ages 60+)	43	15.0%	11.9%
Resources	Poverty (% of adults ages 65+)	37	10.7%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	40	50.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	50	\$17	\$62
and .	High-speed Internet (% of households with adults ages 65+)	31	81.3%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	3	7.9%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	26	51	_
	Volunteerism (% of adults ages 65+)	28	23.3%	22.19
Physical Enviro	onment*	15	0.447	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	25	7.6	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	16	26.1%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	21	27.6%	32.3
Clinical Care*		28	-0.018	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	37	3.7%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	26	30.7	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	39	31.5	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	20	78.3%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	36	65.0%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	25	70.9%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	13	96.8%	96.1
Care	Hospice Care (% of Medicare decedents)	15	50.1%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	24	35.3%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	25	1,389	1,48
Behaviors*		24	0.050	
Nutrition	Exercise (% of adults ages 65+)	23	23.4%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	16	7.6%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	27	31.3%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	26	25.4%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	30	9.1%	8.9%
Health Outcon	nes*	42	-0.442	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	28	7.7%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	20	9.5	9.9
	Excessive Drinking (% of adults ages 65+)	24	7.0%	7.09
	Frequent Mental Distress (% of adults ages 65+)	42	9.7%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	22	17.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	40	2,489	2,15
•	Early Death Racial Disparity (ratio)***	_	1.4	1.5
Physical Health	Falls (% of adults ages 65+)	21	27.7%	27.19
•	Frequent Physical Distress (% of adults ages 65+)	39	16.7%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	34	55%	52%
	Obesity (% of adults ages 65+)	35	32.5%	29.5
	Teeth Extractions (% of adults ages 65+)	36	15.9%	13.49
Overall	,	36	-0.158	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

 Low prevalence of frequent physical distress

State Health Department Website: doh.sd.gov

- · High volunteerism rate
- · High flu vaccination rate

Challenges

- High prevalence of obesity
- Low percentage of households with high-speed internet
- Low home health care worker rate

Highlights

Poverty

43%

from 7.7% to 11.0% of adults ages 65+ between 2019 and 2021.

Physical Inactivity

35%

from 24.6% to 33.1% of adults ages 65+ in fair or better health between 2016 and 2021.

Number of Geriatric Providers

18%

from 14.6 to 17.3 per 100,000 adults ages 65+ between September 2021 and September 2022.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	29	0.070	
	Firearm Deaths (deaths per 100,000 adults ages 65+)+	24	14.6	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	45	17.8	11.8
Economic	Food Insecurity (% of adults ages 60+)	20	10.0%	11.9%
Resources	Poverty (% of adults ages 65+)	38	11.0%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	35	56.9	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	10	\$85	\$62
and	High-speed Internet (% of households with adults ages 65+)	40	79.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	41	18.7%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	13	40	_
	Volunteerism (% of adults ages 65+)	2	39.6%	22.19
Physical Enviro		3	0.960	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	7	5.9	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	9	25.3%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	9	24.8%	32.39
Clinical Care*		26	0.065	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	_	_	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	49	17.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	49	20.8	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	9	79.7%	75.99
Clinical Services	Flu Vaccination (% of adults ages 65+)	2	75.0%	67.59
	Pneumonia Vaccination (% of adults ages 65+)	13	74.2%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	42	93.5%	96.19
Care	Hospice Care (% of Medicare decedents)	45	40.7%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	13	40.1%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	24	1,388	1,48
Behaviors*		19	0.156	
Nutrition	Exercise (% of adults ages 65+)	35	20.5%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	33	5.8%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	32	33.1%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	1	18.2%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	14	8.1%	8.9%
Health Outcon	nes*	25	0.071	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	4	6.0%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	_	_	9.9
	Excessive Drinking (% of adults ages 65+)	31	7.5%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	16	7.4%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	27	18.0	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	15	1,895	2,15
	Early Death Racial Disparity (ratio)***		2.9	1.5
Physical Health	Falls (% of adults ages 65+)	48	33.2%	27.19
	Frequent Physical Distress (% of adults ages 65+)	1	10.6%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	13	46%	52%
	Obesity (% of adults ages 65+)	42	34.3%	29.59
	Teeth Extractions (% of adults ages 65+)	30	13.8%	13.49

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.
Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.
Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of excessive drinking
- High percentage of older adults with a dedicated health care provider
- Low prevalence of severe housing problems

Challenges

- High prevalence of frequent mental distress
- · High prevalence of smoking
- High prevalence of physical inactivity

Highlights

Drug Deaths

26%

from 8.4 to 10.6 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Early Deaths

25%

from 2,241 to 2,793 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

High-speed Internet

23%

from 64.5% to 79.2% of households with adults ages 65+ between 2016 and 2021.

<i>Measure</i>	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	40	-0.350	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	40	19.3	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	40	16.3	11.8
conomic	Food Insecurity (% of adults ages 60+)	37	13.6%	11.9%
Resources	Poverty (% of adults ages 65+)	33	10.2%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	27	64.6	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	44	\$26	\$62
ind	High-speed Internet (% of households with adults ages 65+)	42	79.2%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	4	8.6%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	41	70	_
	Volunteerism (% of adults ages 65+)	41	18.4%	22.1%
Physical Enviro	onment*	11	0.507	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	25	7.6	7.8
Quality	Drinking Water Violations (% of community water systems)	34	0.5%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)*	8	25.2%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	11	25.4%	32.3%
Clinical Care*		36	-0.213	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	31	3.3%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	32	29.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	45	24.5	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	21	78.2%	75.9%
Services	Flu Vaccination (% of adults ages 65+)	31	66.4%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	34	69.9%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	7	97.1%	96.1%
Care	Hospice Care (% of Medicare decedents)	36	45.0%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	41	27.5%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	36	1,612	1,482
Behaviors*		43	-0.770	
Nutrition	Exercise (% of adults ages 65+)	33	20.9%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	34	5.7%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	44	38.5%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	15	23.3%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	49	14.8%	8.9%
Health Outcon	nes*	44	-0.650	
3ehavioral	Cognitive Difficulty (% of adults ages 65+)‡	41	9.2%	7.8%
Behavioral	Drug Deaths (deaths per 100,000 adults ages 65+)*	34	10.6	9.9
Health			2 20/	7.0%
realth	Excessive Drinking (% of adults ages 65+)	1	3.3%	
neaith	Frequent Mental Distress (% of adults ages 65+)	1 47	11.9%	
neaith				8.5%
	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	47	11.9% 21.1 2,793	8.5% 16.9 2,151
N ortality	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)***	47 38 44 —	11.9% 21.1 2,793 1.2	8.5% 16.9 2,151 1.5
Mortality	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+)	47 38 44 — 35	11.9% 21.1 2,793 1.2 29.5%	8.5% 16.9 2,151 1.5 27.1%
Mortality Physical Health	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	47 38 44 — 35 42	11.9% 21.1 2,793 1.2 29.5% 18.4%	8.5% 16.9 2,151 1.5 27.1% 15.8%
Mortality	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	47 38 44 — 35 42 40	11.9% 21.1 2,793 1.2 29.5% 18.4% 56%	8.5% 16.9 2,151 1.5 27.1% 15.8% 52%
Mortality	Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	47 38 44 — 35 42	11.9% 21.1 2,793 1.2 29.5% 18.4%	8.5% 16.9 2,151 1.5 27.1%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.

Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.



Overall Rank

Summary

Strengths

- · High home health care worker rate
- · High prevalence of fruit and vegetable consumption
- Low prevalence of smoking

Challenges

- · High poverty rate
- · Low level of community support expenditures
- · High prevalence of avoiding care due to cost

Highlights

Drug Deaths

from 4.0 to 7.9 deaths per 100,000 adults ages 65+ between 2012-2014 and 2019-2021.

Early Deaths

from 1,857 to 2,439 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

High-speed Internet

from 70.9% to 82.8% of households with adults ages 65+ between 2016 and 2021.

Measure	2S	State Rank	State Value	U.S. Valu
Social & Econo	omic Factors*	42	-0.552	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	27	14.9	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	30	13.8	11.8
Economic	Food Insecurity (% of adults ages 60+)	45	16.3%	11.9%
Resources	Poverty (% of adults ages 65+)	46	11.8%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	25	68.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	46	\$25	\$62
and .	High-speed Internet (% of households with adults ages 65+)	25	82.8%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	36	17.2%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	31	58	_
	Volunteerism (% of adults ages 65+)	35	20.6%	22.1%
Physical Enviro	onment*	39	-0.113	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	44	8.9	7.8
Quality	Drinking Water Violations (% of community water systems)	38	0.8%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	34	31.3%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	31	30.4%	32.39
Clinical Care*	addito agos 62-7	37	-0.274	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	47	5.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	11	41.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	6	79.1	60.3
Preventive Clinical Services	Cancer Screenings (% of adults ages 65-75)	27	77.3%	75.99
	Flu Vaccination (% of adults ages 65+)	33	66.0%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	38	69.0%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	40	94.4%	96.19
Care	Hospice Care (% of Medicare decedents)	16	49.8%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	46	24.1%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	31	1,583	1,482
Behaviors*	,	23	0.098	
Nutrition	Exercise (% of adults ages 65+)	22	23.6%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	6	9.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	36	34.2%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	36	26.8%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	16	8.2%	8.9%
Health Outcon	nes*	30	-0.145	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	32	8.2%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	15	7.9	9.9
	Excessive Drinking (% of adults ages 65+)	28	7.2%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	34	8.6%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	21	16.8	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	38	2,439	2,15
	Early Death Racial Disparity (ratio)**‡	_	1.3	1.5
Physical Health	Falls (% of adults ages 65+)	25	28.3%	27.19
	Frequent Physical Distress (% of adults ages 65+)	23	14.9%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	28	53%	52%
	Obesity (% of adults ages 65+)	23	30.6%	29.5%
	Teeth Extractions (% of adults ages 65+)	28	13.3%	13.49
	-			

 $^{^{\}star}$ Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



State State

115

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Overall Rank

Summary

Strengths

- · Low poverty rate
- · High volunteerism rate
- · High hospice care use

Challenges

- · High suicide rate
- · Low geriatric provider rate
- Low percentage of older adults with a dedicated health care provider

Highlights

Drug Deaths

from 6.9 to 12.4 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Food Insecurity

from 13.7% to 7.4% of adults ages 60+ between 2014-2015 and 2019-2020.

Frequent Mental Distress

from 6.2% to 8.8% of adults ages 65+ between 2017 and 2021.

Measure	28	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	1	0.983	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	26	14.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	39	16.2	11.8
Economic	Food Insecurity (% of adults ages 60+)	4	7.4%	11.9%
Resources	Poverty (% of adults ages 65+)	4	7.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	48	39.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	17	\$53	\$62
and	High-speed Internet (% of households with adults ages 65+)	1	88.6%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	8	10.1%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	1	1	_
	Volunteerism (% of adults ages 65+)	1	44.2%	22.1%
Physical Enviro	onment*	19	0.393	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	33	8.2	7.8
Quality	Drinking Water Violations (% of community water systems)	1	0.0%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)*	17	26.8%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	14	26.4%	32.3%
Clinical Care*		9	0.494	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	18	2.7%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	39	24.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	32	39.7	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	34	75.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	19	69.9%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	14	73.4%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	35	95.4%	96.1%
Care	Hospice Care (% of Medicare decedents)	1	60.7%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	5	45.6%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	4	841	1,482
Behaviors*		3	1.020	
Nutrition	Exercise (% of adults ages 65+)	12	27.0%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	16	7.6%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	4	24.6%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	11	23.0%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	1	4.0%	8.9%
Health Outcon	nes*	9	0.420	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	17	7.0%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	40	12.4	9.9
	Excessive Drinking (% of adults ages 65+)	2	3.9%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	36	8.8%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	32	19.1	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	10	1,753	2,151
	Early Death Racial Disparity (ratio)**‡	_	2.0	1.5
Physical Health	Falls (% of adults ages 65+)	36	29.8%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	19	14.6%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	16	47%	52%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value)

Obesity (% of adults ages 65+)

Teeth Extractions (% of adults ages 65+)

Overall

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



18

28.7%

9.7%

0.722

29.5%

13.4%

Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of multiple chronic conditions
- High volunteerism rate
- · Low prevalence of physical inactivity

Challenges

- · High suicide rate
- · High prevalence of excessive drinking
- Low hospice care use

Highlights

Poverty

from 6.1% to 9.1% of adults ages 65+ between 2019 and 2021.

Depression

from 14.2% to 19.7% of adults ages 65+ between 2018 and 2021.

Flu Vaccination Rates

from 64.2% to 73.5% of adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	3	0.841	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	28	15.2	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	22	11.7	11.8
Economic	Food Insecurity (% of adults ages 60+)	15	9.6%	11.9%
Resources	Poverty (% of adults ages 65+)	16	9.1%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	6	\$120	\$62
and	High-speed Internet (% of households with adults ages 65+)	20	83.5%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	15	11.7%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	11	37	_
	Volunteerism (% of adults ages 65+)	4	36.0%	22.1%
Physical Enviro	onment*	13	0.500	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	4	5.0	7.8
Quality	Drinking Water Violations (% of community water systems)	30	0.3%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	36	32.2%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	40	35.0%	32.3%
Clinical Care*	,	17	0.226	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	11	2.4%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	23	33.8	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	30	40.3	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	12	79.2%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	6	73.5%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	29	70.6%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	27	96.1%	96.1%
Care	Hospice Care (% of Medicare decedents)	37	44.5%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	37	29.3%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	19	1,271	1,482
Behaviors*	,	1	1.058	
Nutrition	Exercise (% of adults ages 65+)	3	29.3%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	1	13.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or	4	24.6%	31.2%
	better health)	4.0	00.111	00.00
Sleep Health	Insufficient Sleep (% of adults ages 65+)	12	23.1%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	32	9.2%	8.9%
Health Outcon	nes*	16	0.277	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	14	6.8%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	14	7.5	9.9
	Excessive Drinking (% of adults ages 65+)	44	8.9%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	16	7.4%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	40	22.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	4	1,643	2,151
	Early Death Racial Disparity (ratio)***			1.5
Physical Health	Falls (% of adults ages 65+)	45	31.1%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	8	12.5%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	3	38%	52%
	Obesity (% of adults ages 65+)	12	26.8%	29.5%
	Teeth Extractions (% of adults ages 65+)	37	16.0%	13.4%
Overall		5	0.603	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

Non-ranking measure.

— Data not available, missing or suppressed.

Virginia

State Health Department Website: vdh.virginia.gov

Overall Rank

Summary

Strengths

- Low prevalence of frequent mental distress
- · Low poverty rate
- · High pneumonia vaccination rate

Challenges

- High prevalence of obesity
- · Low level of community support expenditures
- · Low home health care worker rate

Highlights

Drug Deaths

from 3.6 to 7.0 deaths per 100,000 adults ages 65+ between 2014-2016 and 2019-2021.

Number of Geriatric Providers

from 25.5 to 31.8 per 100,000 adults ages 65+ between September 2019 and September 2022.

Early Deaths

from 1,736 to 2,105 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Measure	28	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	18	0.299	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	22	14.4	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	20	11.5	11.8
Economic	Food Insecurity (% of adults ages 60+)	20	10.0%	11.9%
Resources	Poverty (% of adults ages 65+)	6	8.0%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	26	67.4	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	37	\$31	\$62
and	High-speed Internet (% of households with adults ages 65+)	21	83.3%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	19	12.9%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	17	45	_
	Volunteerism (% of adults ages 65+)	26	23.9%	22.1%
Physical Enviro		10	0.537	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	12	6.6	7.8
Quality	Drinking Water Violations (% of community water systems)	14	0.1%	0.8%
Housing	Housing Cost Burden (% of households with adults	26	28.6%	31.8%
	ages 65+)‡			
	Severe Housing Problems (% of small households with adults ages 62+)	26	29.4%	32.3%
Clinical Care*		25	0.104	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	27	3.1%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	25	31.8	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	33	39.2	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	11	79.3%	75.9%
Clinical Services	Flu Vaccination (% of adults ages 65+)	24	69.2%	67.5%
	Pneumonia Vaccination (% of adults ages 65+)	9	75.0%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	25	96.2%	96.1%
Care	Hospice Care (% of Medicare decedents)	30	46.2%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	45	24.2%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	23	1,382	1,482
Behaviors*		16	0.206	
Nutrition	Exercise (% of adults ages 65+)	27	21.8%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	27	6.6%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	17	28.7%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	15	23.3%	26.0%
Tobacco Use	Smoking (% of adults ages 65+)	14	8.1%	8.9%
Health Outcon	mes*	18	0.272	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	20	7.2%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	9	7.0	9.9
	Excessive Drinking (% of adults ages 65+)	12	6.0%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	6	7.0%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	22	17.2	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	27	2,105	2,151
,	Early Death Racial Disparity (ratio)***		1.4	1.5
Physical Health	Falls (% of adults ages 65+)	11	25.9%	27.1%
, 0.00. 1100101	Frequent Physical Distress (% of adults ages 65+)	19	14.6%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	26	52%	52%
		31	31.6%	29.5%
	Obesity (% of adults ages 65+) Teeth Extractions (% of adults ages 65+)	31 11	31.6% 10.9%	29.5% 13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Washington

Overall Rank

Summary

Strengths

- · Low prevalence of multiple chronic conditions
- · Low poverty rate
- · Low prevalence of physical inactivity

Challenges

- · High suicide rate
- Low level of community support expenditures
- · Low hospice care use

Highlights

Drug Deaths

from 6.0 to 11.7 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Volunteerism

from 36.8% to 25.6% of adults ages 65+ between 2017 and 2021.

High-speed Internet

from 78.9% to 88.1% of households with adults ages 65+ between 2016 and 2021.

Family Safety Economic Resources Social Support and Engagement	Firearm Deaths (deaths per 100,000 adults ages 65+)* Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)* Food Insecurity (% of adults ages 60+) Poverty (% of adults ages 65+)	5 22 4 6	0.698 14.4 7.3	13.1
Family Safety Economic Resources Social Support and Engagement	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡ Food Insecurity (% of adults ages 60+)	6	7.3	
Economic Resources Social Support and Engagement	Food Insecurity (% of adults ages 60+)	6		1
Resources Social Support and Engagement				11.8
Social Support and Engagement	Poverty (% of adults ages 65+)	_	8.4%	11.99
and Engagement		7	8.2%	10.39
and Engagement	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
and Engagement	Community Support Expenditures (dollars per adult ages 60+)	47	\$24	\$62
	High-speed Internet (% of households with adults ages 65+)	2	88.1%	83.19
F	Low-care Nursing Home Residents (% of residents)	9	10.5%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	14	42	_
	Volunteerism (% of adults ages 65+)	20	25.6%	22.19
Physical Enviro	nment*	38	-0.093	
	Air Pollution (micrograms of fine particles per cubic meter)	32	8.0	7.8
Quality	Drinking Water Violations (% of community water systems)	34	0.5%	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	40	32.9%	31.8
	Severe Housing Problems (% of small households with adults ages 62+)	37	33.9%	32.3
Clinical Care*		8	0.539	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	14	2.6%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	22	34.0	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	14	50.1	60.
Preventive	Cancer Screenings (% of adults ages 65-75)	14	79.0%	75.9
Clinical Services	Flu Vaccination (% of adults ages 65+)	15	71.5%	67.5
	Pneumonia Vaccination (% of adults ages 65+)	3	75.4%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	25	96.2%	96.1
Care	Hospice Care (% of Medicare decedents)	33	45.5%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	14	39.7%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	6	930	1,48
Behaviors*	,	5	0.862	
Nutrition	Exercise (% of adults ages 65+)	7	27.9%	23.1
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	7	9.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or	3	22.2%	31.2
Sleep Health	better health)	22	25.1%	26.0
•	Insufficient Sleep (% of adults ages 65+) Smoking (% of adults ages 65+)	12	7.8%	8.99
		14		0.57
Health Outcom			0.340	
	Cognitive Difficulty (% of adults ages 65+)+	33	8.3%	7.89
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	38	11.7	9.9
	Excessive Drinking (% of adults ages 65+)	29	7.3%	7.09
	Frequent Mental Distress (% of adults ages 65+)	29	8.3%	8.59
	Suicide (deaths per 100,000 adults ages 65+)	35	19.9	16.9
	Early Death (deaths per 100,000 adults ages 65-74)	14	1,855	2,15
	Early Death Racial Disparity (ratio)***		2.1	1.5
•	Falls (% of adults ages 65+)	41	30.6%	27.19
	Frequent Physical Distress (% of adults ages 65+)	22	14.8%	15.8
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	6	41%	529
	Obesity (% of adults ages 65+) Teeth Extractions (% of adults ages 65+)	13 8	26.9% 9.9%	29.5 13.4

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

West Virginia State Health Department Website: dhhr.wv.gov Overall Rank 47

Summary

Strengths

- Low prevalence of excessive drinking
- High level of community support expenditures
- Low prevalence of severe housing problems

Challenges

- High prevalence of frequent mental distress
- High prevalence of multiple chronic conditions
- High prevalence of food insecurity

Highlights

Physical Inactivity

34%

from 30.5% to 41.0% of adults ages 65+ in fair or better health between 2016 and 2021.

High-speed Internet

24%

from 60.4% to 75.0% of households with adults ages 65+ between 2016 and 2021.

Early Deaths

18%

from 2,589 to 3,066 deaths per 100,000 adults ages 65-74 between 2020 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	43	-0.648	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	42	19.8	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	33	14.7	11.8
Economic Resources	Food Insecurity (% of adults ages 60+)	46	16.9%	11.9%
	Poverty (% of adults ages 65+)	43	11.5%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	1	100.0	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	13	\$68	\$62
nd	High-speed Internet (% of households with adults ages 65+)	49	75.0%	83.1%
ngagement	Low-care Nursing Home Residents (% of residents)	38	17.5%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	45	82	_
	Volunteerism (% of adults ages 65+)	33	22.1%	22.1%
Physical Enviro		31	0.107	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	19	7.3	7.8
Quality	Drinking Water Violations (% of community water systems)	50	10.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	1	19.1%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	1	18.6%	32.3%
Clinical Care*		44	-0.604	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	38	3.8%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	42	21.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	25	42.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	35	75.0%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	41	63.7%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	20	72.0%	69.79
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	12	96.9%	96.1%
Care	Hospice Care (% of Medicare decedents)	40	42.4%	46.7%
	Nursing Home Quality (% of beds rated four or five stars)	47	22.5%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	50	2,568	1,482
Behaviors*		50	-1.530	
Nutrition	Exercise (% of adults ages 65+)	47	13.5%	23.1%
ınd Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	47	4.0%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	48	41.0%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	47	30.1%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	40	11.4%	8.9%
Health Outcon	nes*	50	-1.304	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	46	9.6%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	45	14.1	9.9
	Excessive Drinking (% of adults ages 65+)	5	4.6%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	49	12.6%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	39	21.3	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	49	3,066	2,151
-	Early Death Racial Disparity (ratio)***	_	1.1	1.5
hysical Health	Falls (% of adults ages 65+)	46	32.4%	27.1%
-	Frequent Physical Distress (% of adults ages 65+)	49	22.4%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	48	60%	52%
	Obesity (% of adults ages 65+)	49	35.9%	29.5%
	Teeth Extractions (% of adults ages 65+)	49	22.2%	13.4%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value). Higher summation scores are better. Scores are not calculated for the District of Columbia.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.
Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value.
Ties in value are assigned equal ranks.



^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

Strengths

- Low prevalence of frequent mental distress
- · Low risk of social isolation
- Low prevalence of physical inactivity

Challenges

- High prevalence of excessive drinking
- Low percentage of households with high-speed internet
- · Low geriatric provider rate

Highlights

Drug Deaths

113%

from 3.9 to 8.3 deaths per 100,000 adults ages 65+ between 2011-2013 and 2019-2021.

Flu Vaccination Rates

39%

from 52.7% to 73.2% of adults ages 65+ between 2017 and 2021.

Poverty

18%

from 7.4% to 8.7% of adults ages 65+ between 2019 and 2021.

Measure	es es	State Rank	State Value	U.S. Valu
Social & Economic Factors*		15	0.355	
	Firearm Deaths (deaths per 100,000 adults ages 65+)+	16	12.3	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	12	10.2	11.8
Economic Resources	Food Insecurity (% of adults ages 60+)	12	9.5%	11.9%
	Poverty (% of adults ages 65+)	13	8.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	18	84.3	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	21	\$48	\$62
and Engagement	High-speed Internet (% of households with adults ages 65+)	33	80.8%	83.19
	Low-care Nursing Home Residents (% of residents)	12	11.0%	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	8	35	_
	Volunteerism (% of adults ages 65+)	30	22.7%	22.19
Physical Enviro		48	-0.517	
Air and Water Quality	Air Pollution (micrograms of fine particles per cubic meter)	31	7.9	7.8
	Drinking Water Violations (% of community water systems)	46	4.4%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	31	29.7%	31.89
	Severe Housing Problems (% of small households with adults ages 62+)	34	31.5%	32.3
Clinical Care*		12	0.412	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	7	2.3%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	31	29.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	8	69.0	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	47	68.3%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	7	73.2%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	22	71.8%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	14	96.7%	96.19
Care	Hospice Care (% of Medicare decedents)	4	54.7%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	9	42.8%	32.69
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	17	1,268	1,48
Behaviors*	•	6	0.754	
Nutrition	Exercise (% of adults ages 65+)	1	30.3%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	29	6.3%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	8	26.4%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	7	22.0%	26.0
Tobacco Use	Smoking (% of adults ages 65+)	9	7.6%	8.9%
Health Outcon	nes*	17	0.273	
Behavioral	Cognitive Difficulty (% of adults ages 65+)*	1	5.3%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	16	8.3	9.9
	Excessive Drinking (% of adults ages 65+)	47	10.3%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	3	6.0%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	15	15.7	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	13	1,848	2,15
•	Early Death Racial Disparity (ratio)***	_	2.1	1.5
Physical Health	Falls (% of adults ages 65+)	16	26.9%	27.19
	Frequent Physical Distress (% of adults ages 65+)	13	13.5%	15.89
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	13	46%	52%
	Obesity (% of adults ages 65+)	33	32.3%	29.5
	Teeth Extractions (% of adults ages 65+)	13	11.2%	13.49
Overall		11	270	.5. 17

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia.

^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org.

Strengths

- · Low prevalence of multiple chronic conditions
- · Low poverty rate
- · High level of community support expenditures

Challenges

- · High suicide rate
- · Low hospice care use
- · Low percentage of older adults with a dedicated health care provider

Highlights

Early Deaths

from 1,720 to 2,299 deaths per 100,000 adults ages 65-74 between 2019 and 2021.

Number of Geriatric Providers

from 14.7 to 19.3 per 100,000 adults ages 65+ between September 2019 and September 2022.

Volunteerism

from 32.6% to 39.2% of adults ages 65+ between 2017 and 2021.

Measure	es es	State Rank	State Value	U.S. Value
Social & Econo	omic Factors*	8	0.536	
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	50	34.5	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)*	50	24.1	11.8
Economic Resources	Food Insecurity (% of adults ages 60+)	31	12.6%	11.9%
	Poverty (% of adults ages 65+)	1	6.7%	10.3%
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	50	17.7	81.0
Social Support	Community Support Expenditures (dollars per adult ages 60+)	4	\$139	\$62
ind	High-speed Internet (% of households with adults ages 65+)	17	85.0%	83.1%
Engagement	Low-care Nursing Home Residents (% of residents)	43	21.4%	15.2%
	Risk of Social Isolation (index from 1-100, adults ages 65+)	8	35	_
	Volunteerism (% of adults ages 65+)	3	39.2%	22.1%
Physical Enviro	onment*	1	1.310	
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	3	4.7	7.8
Quality	Drinking Water Violations (% of community water systems)	24	0.2%	0.8%
Housing	Housing Cost Burden (% of households with adults ages 65+)‡	2	23.9%	31.8%
	Severe Housing Problems (% of small households with adults ages 62+)	2	22.9%	32.3%
Clinical Care*	,	50	-0.976	
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	28	3.2%	3.4%
	Geriatric Providers (providers per 100,000 adults ages 65+)	48	19.3	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	36	33.4	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	50	62.4%	75.9%
Clinical	Flu Vaccination (% of adults ages 65+)	47	60.5%	67.5%
Services	Pneumonia Vaccination (% of adults ages 65+)	41	66.8%	69.7%
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	48	91.3%	96.1%
Care	Hospice Care (% of Medicare decedents)	47	33.1%	46.79
	Nursing Home Quality (% of beds rated four or five stars)	29	32.8%	32.6%
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	11	1,137	1,482
Behaviors*		14	0.308	
Nutrition	Exercise (% of adults ages 65+)	17	25.0%	23.1%
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	25	6.9%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	19	29.3%	31.2%
Sleep Health	Insufficient Sleep (% of adults ages 65+)	8	22.5%	26.09
Tobacco Use	Smoking (% of adults ages 65+)	28	9.0%	8.9%
Health Outcon	nes*	33	-0.177	
Behavioral	Cognitive Difficulty (% of adults ages 65+)‡	33	8.3%	7.8%
Health	Drug Deaths (deaths per 100,000 adults ages 65+)*	31	10.4	9.9
	Excessive Drinking (% of adults ages 65+)	33	7.7%	7.0%
	Frequent Mental Distress (% of adults ages 65+)	19	7.6%	8.5%
	Suicide (deaths per 100,000 adults ages 65+)	50	38.8	16.9
Mortality	Early Death (deaths per 100,000 adults ages 65-74)	34	2,299	2,151
	Early Death Racial Disparity (ratio)***		2.5	1.5
Physical Health	Falls (% of adults ages 65+)	44	30.7%	27.1%
	Frequent Physical Distress (% of adults ages 65+)	40	16.9%	15.8%
	Multiple Chronic Conditions (% of Medicare beneficiaries ages 65-74)	1	35%	52%
	Obesity (% of adults ages 65+)	21	29.6%	29.5%
	Teeth Extractions (% of adults ages 65+)	37	16.0%	13.4%
Overall		26	0.048	

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks.



Higher summation scores are better. Scores are not calculated for the District of Columbia

 $^{^{\}star\star}$ Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.

District of Columbia

Summary

Strengths

- · Low suicide rate
- · High level of community support expenditures
- · High geriatric provider rate

Challenges

- · High prevalence of excessive drinking
- · High poverty rate
- · Low hospice care use

Highlights

Drug Deaths

from 39.6 to 75.3 deaths per 100,000 adults ages 65+ between 2016-2018 and 2019-2021.

Flu Vaccination Rates

from 51.7% to 70.8% of adults ages 65+ between 2016 and 2021.

High-speed Internet

from 68.5% to 80.0% of households with adults ages 65+ between 2017 and 2021.

Measure	2S	D.C. Value	U.S Valu
Social & Econo	omic Factors*		
Community and	Firearm Deaths (deaths per 100,000 adults ages 65+)‡	_	13.1
Family Safety	Motor Vehicle Deaths (deaths per 100,000 adults ages 65+)‡	3.3	11.8
Economic	Food Insecurity (% of adults ages 60+)	18.8%	11.99
Resources	Poverty (% of adults ages 65+)	13.8%	10.39
	SNAP Reach (participants per 100 adults ages 60+ in poverty)	93.4	81.C
Social Support	Community Support Expenditures (dollars per adult ages 60+)	\$338	\$62
and .	High-speed Internet (% of households with adults ages 65+)	80.0%	83.19
Engagement	Low-care Nursing Home Residents (% of residents)	_	15.29
	Risk of Social Isolation (index from 1-100, adults ages 65+)	_	_
	Volunteerism (% of adults ages 65+)	31.2%	22.19
Physical Enviro			
Air and Water	Air Pollution (micrograms of fine particles per cubic meter)	8.7	7.8
Quality	Drinking Water Violations (% of community water systems)	_	0.89
Housing	Housing Cost Burden (% of households with adults ages 65+)*	39.4%	31.89
ŭ	Severe Housing Problems (% of small households with adults	39.0%	32.3
	ages 62+)		
Clinical Care*			
Access to Care	Avoided Care Due to Cost (% of adults ages 65+)	2.9%	3.49
	Geriatric Providers (providers per 100,000 adults ages 65+)	90.9	36.4
	Home Health Care Workers (workers per 1,000 adults ages 65+)	127.1	60.3
Preventive	Cancer Screenings (% of adults ages 65-75)	83.2%	75.9
Clinical	Flu Vaccination (% of adults ages 65+)	70.8%	67.5
Services	Pneumonia Vaccination (% of adults ages 65+)	72.2%	69.7
Quality of	Dedicated Health Care Provider (% of adults ages 65+)	95.6%	96.1
Care	Hospice Care (% of Medicare decedents)	25.8%	46.7
	Nursing Home Quality (% of beds rated four or five stars)	31.5%	32.6
	Preventable Hospitalizations (discharges per 100,000 Medicare beneficiaries ages 65-74)	2,273	1,48
Behaviors*			
Nutrition	Exercise (% of adults ages 65+)	29.8%	23.19
and Physical	Fruit and Vegetable Consumption (% of adults ages 65+)	13.9%	7.3%
Activity	Physical Inactivity (% of adults ages 65+ in fair or better health)	26.7%	31.29
Sleep Health	Insufficient Sleep (% of adults ages 65+)	26.9%	26.0
	Smoking (% of adults ages 65+)	10.6%	8.99
Tobacco Use			
Tobacco Use Health Outco n			
	nes*	8.2%	7.89
Health Outcon		8.2% 75.3	
Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)*	75.3	9.9
Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+)	75.3 9.5%	7.8% 9.9 7.0% 8.5%
Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+)	75.3 9.5% 7.0%	9.9 7.0% 8.5%
Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+)	75.3 9.5% 7.0% 7.6	9.9 7.0% 8.5% 16.9
Health Outcon Behavioral	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74)	75.3 9.5% 7.0% 7.6 2,595	9.9 7.0% 8.5% 16.9 2,15
Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡	75.3 9.5% 7.0% 7.6 2,595 4.1	9.9 7.0% 8.5% 16.9 2,15 1.5
Health Outcon Behavioral Health	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+)	75.3 9.5% 7.0% 7.6 2,595 4.1 29.1%	9.9 7.0% 8.5% 16.9 2,15 1.5
Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)* Drug Deaths (deaths per 100,000 adults ages 65+)* Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)*** Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+) Multiple Chronic Conditions (% of Medicare beneficiaries	75.3 9.5% 7.0% 7.6 2,595 4.1	9.9 7.09 8.59 16.9 2,15 1.5 27.19
Health Outcon Behavioral Health Mortality	Cognitive Difficulty (% of adults ages 65+)‡ Drug Deaths (deaths per 100,000 adults ages 65+)‡ Excessive Drinking (% of adults ages 65+) Frequent Mental Distress (% of adults ages 65+) Suicide (deaths per 100,000 adults ages 65+) Early Death (deaths per 100,000 adults ages 65-74) Early Death Racial Disparity (ratio)**‡ Falls (% of adults ages 65+) Frequent Physical Distress (% of adults ages 65+)	75.3 9.5% 7.0% 7.6 2,595 4.1 29.1% 11.6%	9.9 7.0% 8.5%

^{*} Value is a sum of weighted, ranking measure z-scores (the number of standard deviations a state value was above or below the U.S. value).

For measure descriptions, source details and methodology, visit AmericasHealthRankings.org. Ranks are calculated by state for each measure, ordered by the states' values, with 1 corresponding to the healthiest value and 50 the least healthy value. Ties in value are assigned equal ranks



Higher summation scores are better. Scores are not calculated for the District of Columbia

^{**} Disparity measures compare the group with the highest or lowest rate to the white rate.

^{*} Non-ranking measure.

Data not available, missing or suppressed.